arXiv:hep-ph/0107272v3 22 Mar 2002

Scalar perturbations during multiple-field slow-roll inflation

S Groot Nibbelink† and B J W van Tent‡

†Physikalisches Institut, Universität Bonn, Nußallee 12, D-53115 Bonn, Germany
‡Spinoza Institute/Institute for Theoretical Physics, Utrecht University, P.O. Box 80195, 3508 TD Utrecht, The Netherlands

E-mail: nibblink@th.physik.uni-bonn.de, B.J.W.vanTent@phys.uu.nl

Abstract. We calculate the scalar gravitational and matter perturbations in the context of slow-roll inflation with multiple scalar fields, that take values on a (curved) manifold, to first order in slow roll. For that purpose a basis for these perturbations determined by the background dynamics is introduced and multiple-field slow-roll functions are defined. To obtain analytic solutions to first order, the scalar perturbation modes have to be treated in three different regimes. Matching is performed by analytically identifying leading order asymptotic expansions in different regions. Possible sources for multiple-field effects in the gravitational potential are the particular solution caused by the coupling to the field perturbation perpendicular to the field velocity, and the rotation of the basis. The former can contribute even to leading order if the corresponding multiple-field slow-roll function is sizable during the last 60 e-folds. Making some simplifying assumptions, the evolution of adiabatic and isocurvature perturbations after inflation is discussed. The analytical results are illustrated and checked numerically with the example of a quadratic potential.

PACS number: 98.80.Cq

1. Introduction

As has been known for a long time, inflation [1, 2] offers a mechanism for the production of density perturbations, which are supposed to be the seeds for the formation of large scale structures in the universe. This mechanism is the magnification of microscopic quantum fluctuations in the scalar fields present during the inflationary epoch into macroscopic matter and metric perturbations. Also, since a part of the primordial spectrum of density perturbations is observed in the cosmic microwave background radiation (CMBR), this mechanism offers one of the most important ways of checking and constraining possible models of inflation, see e.g. [3], especially when combined with large scale structure data [4].

The theory of the production of density perturbations in inflation has been studied for a long time. First in the case of a single real scalar field (see e.g. [5, 6, 7, 8, 9, 10, 11]), but later also for multiple fields, for example in the following papers. Pioneering work was done in [12, 13]. Using gauge invariant variables the authors of [14, 15, 16, 17, 18] treated two-field inflation models. The fluid flow approach was extended to multiple fields in [19], while a more geometrical approach was used in [20, 21]; both methods assumed several slow-roll-like conditions on the potential. Using slow-roll approximations for both the background and the perturbation equations the authors of [22, 23, 24] were able to find expressions for the metric perturbations in multiple-field inflation. The authors of [25, 26] paid special attention to gauge issues in their discussion of multiple-field perturbations. The case of perturbations in generalized gravity theories was studied in [27, 28, 29].

There are two important reasons for considering inflation with multiple scalar fields. To realize sufficient inflation before a graceful exit from the inflationary era and produce the observed density perturbation spectrum in a model without very unnatural values of the parameters and initial conditions, one is naturally led to the introduction of additional fields. This is the motivation for hybrid inflation models [30] (related models can be found in [19]). The other reason is that many theories beyond the standard model of particle physics, like grand unification, supersymmetry or effective supergravity from string theory, contain a lot of scalar fields. Ultimately one would hope to be able to identify those fields that can act as inflatons. In addition such string-inspired supersymmetric models naturally have non-minimal kinetic terms due to a Kähler potential that is not of the minimal form $G = \bar{z}z$ for a complex scalar field z (perhaps a modulus of string theory) contained in a chiral multiplet. (Since a complex field can always be written as two real fields, it is sufficient to consider only real scalar fields in our paper.)

To be able to investigate the possibility that such models can produce (sufficient) inflation and density perturbations in accordance with present and future observations, it is necessary to have a general treatment available that can handle an arbitrary number of scalar fields with an arbitrary (Kähler) field metric and a generic potential. Most of the previous literature on multiple-field inflationary density perturbations is limited with respect to these aspects, usually by considering only two fields and minimal kinetic terms. An exception to this are the papers [20, 21], but these still left space for improvement, most importantly regarding the treatment of slow roll, the rotation of background fields, the transition region, and the analysis of the particular solution for the gravitational potential caused by the coupling to multiple fields. In this paper we provide a general treatment by computing the scalar gravitational and matter perturbations to first order in slow roll during inflation with multiple real scalar fields that may have non-minimal kinetic terms. Which of these fields acts as inflaton during which part of the inflationary period is determined automatically in our formalism and does not have to be specified beforehand. The treatment has three important ingredients: a geometrical setup, generalized slow-roll functions, and a thorough discussion of the period when the scales of the perturbation modes are of the same order of magnitude as the Hubble scale. Let us elaborate on these points.

First of all, the scalar fields and their perturbations have to be described in a geometrical way to make certain that their physical description is independent of the coordinate system that parameterizes the field manifold. To describe the evolution and quantization of the perturbations it is essential to introduce a background-field induced basis (of the tangent bundle), such that the components of the perturbations in this basis are canonically normalized so that only physical degrees of freedom are quantized. This basis is also used to distinguish effectively single-field and truly multiple-field effects and to identify adiabatic and entropy perturbations during inflation.

We generalize the slow-roll parameters for a single background field to multiple scalar fields in a systematic way. They are defined independently of the coordinates used on the scalar manifold. In addition, our definition of these variables is such that they apply to any background time variable (e.g. comoving or conformal time). In our paper we need this generality to rigorously define the notion of slow roll applied to the perturbations. Our slow-roll functions are defined independently of whether slow roll is valid, and can be viewed as short-hand notation to denote derivatives of the Hubble parameter and the field velocity. As such they can be identified in all kinds of exact equations, making it clear what the behaviour of these equations is if the system is in the slow-roll regime. The relevance of the slow-roll function $\tilde{\eta}^{\perp}$ should be stressed at this point, as it is the measure of true multiple-field effects. Indeed, by definition $\tilde{\eta}^{\perp}$ vanishes in the case of a single scalar field model. The multiple-field character of inflation is only apparent when $\tilde{\eta}^{\perp}$ is non-negligible. In particular, it determines the size of the multiple-field contributions to the adiabatic perturbation and the mixing between adiabatic and isocurvature perturbations.

During inflation there is a relatively sharp transition in the behaviour of a fluctuation when the corresponding wavelength becomes larger than the Hubble radius ('passes through the horizon'); this moment identifies a certain scale k. For observationally interesting scales from the point of view of inflation (those that reentered the horizon only after the time of recombination when the CMBR was formed) this happened approximately 60 e-folds before the end of inflation [7]. To find results for the perturbations valid to first order in slow roll this transition has to be treated very carefully. We determine the order of various contributions, in particular multiple-field rotation effects, to see if they can be neglected to first order or should be taken into account. Analytic properties of the solutions in the different regions are used to relate the normalization factor at the end of inflation to the initial conditions at a much earlier time before the transition.

The main subject of this paper is the treatment of perturbations during inflation, but we also provide results for adiabatic and isocurvature perturbations and the correlations between them at the time of recombination when the CMBR was formed. Although adiabatic perturbations are always present, this is not true for isocurvature perturbations. Whether the latter survive till recombination at all depends on additional physical assumptions. Here we follow [24] regarding assumptions about the decay of the scalar fields and the evolution after inflation, which are such that isocurvature perturbations do survive in principle. We also neglect the effects of preheating. A more thorough treatment of these aspects will be the subject of another paper.

Apart from this introduction the paper is structured as follows. In section 2.1 the background with multiple scalar fields is described using geometrical concepts which are explained in Appendix A. An orthonormal basis induced by the dynamics of the background fields is also introduced here. Section 2.2 then describes the multiple-field slow-roll formalism.

Section 3 is devoted to the perturbations in multiple-field inflation and is the main part of this paper. In section 3.1 the equations of motion for the scalar gravitational and matter perturbations are derived, and the choice of perturbation variables is discussed. The next section 3.2 focuses on the quantization of the dynamical scalar perturbations. After discussing the outlines of the calculation in section 3.3, and introducing the concept of slow roll on the perturbations in section 3.4, solving the equations to find expressions for the perturbations, in particular the gravitational potential, in terms of background quantities only is done in section 3.5. Section 3.6 then deals with the perturbations after inflation and gives expressions for the vacuum correlators of the gravitational potential at the time of recombination.

In section 4 the example of a quadratic potential with multiple scalar fields is discussed, not only to illustrate the theory of section 3, but also as a numerical check of our analytical results. Analytical expressions for this example are derived in section 4.1, while section 4.2 gives numerical results. The results of this paper are summarized and discussed in section 5.

2. Slow-roll background in multiple-field inflation

2.1. Equations of motion for the background

The background of the universe is described by the flat Robertson-Walker metric in terms of a general time variable τ:

$$\mathrm{d}s^2 = -b^2 \mathrm{d}\tau^2 + a^2 \mathrm{d}\boldsymbol{x}^2 \tag{1}$$

with $a(\tau)$ the spatial scale factor. The temporal scale factor b is defined by the specific choice of time variable: for comoving time t and conformal time η it is given by $b = 1$ and $b = a$, respectively, leading to the relation $\mathrm{d}t = a\mathrm{d}\eta$. The main reason for setting up the background equations in terms of a general time variable is that we need those in our discussion of slow roll on the perturbations in section 3.4. In addition, since different equations are best solved using different time variables, it is convenient to set up the formalism for a general time variable, to avoid repetitions of almost identical equations. Moreover, with this approach various definitions and conclusions are manifestly independent of the choice of time coordinate, in particular those related to the slow-roll approximation. A derivative with respect to the general time variable τ is denoted by $^{;} \equiv \partial_\tau$, one with respect to comoving time by $\dot{} \equiv \partial_t$, and one with respect to conformal time by $' \equiv \partial_\eta$. Hubble parameters $H_a \equiv \partial_\tau a/a$ and

$H_b \equiv \partial_\tau b/b$ are associated with the scale factors a and b. For H_a in terms of comoving and conformal time we define the conventional symbols: $H = \dot{a}/a$ and $\mathcal{H} = a'/a = aH$.

For the matter part of the universe we consider an arbitrary number of real scalar fields that are the components of a vector $\boldsymbol{\phi}$ and the coordinates on a possibly non-trivial field manifold $\mathcal{M}$ with metric $\mathbf{G}$. The Lagrangean for the scalar field theory with a potential V on this manifold in a general spacetime that is quadratic in the derivatives can be written as

$$\mathcal{L}_{\mathcal{M}} = \sqrt{-g}\left(-\frac{1}{2}\partial^\mu\boldsymbol{\phi}\cdot\partial_\mu\boldsymbol{\phi} - V(\boldsymbol{\phi})\right) = \sqrt{-g}\left(-\frac{1}{2}g^{\mu\nu}\partial_\mu\boldsymbol{\phi}^T\mathbf{G}\partial_\nu\boldsymbol{\phi} - V(\boldsymbol{\phi})\right), \tag{2}$$

with g the determinant of $g_{\mu\nu}$. Notice that the kinetic term contains both the inverse spacetime metric $g^{\mu\nu}$ and the field metric $\mathbf{G}$. Definitions of various geometrical concepts like the inner product $\boldsymbol{A}\cdot\boldsymbol{B} = \boldsymbol{A}^\dagger\boldsymbol{B} = \boldsymbol{A}^T\mathbf{G}\boldsymbol{B}$ and the derivatives $\mathcal{D}_\mu$ (with respect to spacetime) and $\boldsymbol{\nabla}$ (with respect to the fields), that are covariant with respect to the geometry of the manifold $\mathcal{M}$, can be found in Appendix A.

This geometrical description of the kinetic part of the Lagrangean is motivated by high-energy theoretical models where non-minimal kinetic terms appear naturally. In particular, in supergravity models [31, 32], in which renormalizability is no longer an issue, supersymmetry forces the scalar Lagrangean to take the form

$$\mathcal{L} = -G_{\bar{\alpha}\alpha}\partial_\mu\bar{z}^{\underline{\alpha}}\partial^\mu z^\alpha - V, \qquad\qquad V = e^{-G}\Big(G_\alpha G^{\alpha\underline{\alpha}}G_{\underline{\alpha}} - 3\Big). \tag{3}$$

Here the subscripts $_\alpha$ and $_{\underline{\alpha}}$ denote differentiation with respect to the complex coordinates z^α and their conjugates $\bar{z}^{\underline{\alpha}}$, respectively. These coordinates parameterize a so-called Kähler manifold, which has the property that the metric $G_{\underline{\alpha}\alpha}$ (with $G^{\alpha\underline{\alpha}}$ its inverse) can be determined from the Kähler potential G as its second mixed derivative. This Kähler potential $G = K + \ln|W|^2$ consists of a Kähler potential $K(z,\bar{z})$ of the complex manifold and a holomorphic superpotential $W(z)$. Models of (heterotic) string theory obtained by compactification lead to supergravity theories with the structure of the scalar field theory described above, see [33]. The resulting Kähler potential K has typical no-scale supergravity features [34] due to the strong requirement of modular invariance [35]. An introduction to string phenomenology can be found in [36].

The equations of motion for the scalars are given by

$$g^{\mu\nu}\left(\mathcal{D}_\mu\delta^\lambda_\nu - \Gamma^\lambda_{\mu\nu}\right)\partial_\lambda\boldsymbol{\phi} - \mathbf{G}^{-1}\boldsymbol{\nabla}^T V = 0, \tag{4}$$

and the Einstein equations read

$$\frac{1}{\kappa^2}G^\mu_\nu = T^\mu_\nu = \partial^\mu\boldsymbol{\phi}\cdot\partial_\nu\boldsymbol{\phi} - \delta^\mu_\nu\left(\frac{1}{2}\partial^\lambda\boldsymbol{\phi}\cdot\partial_\lambda\boldsymbol{\phi} + V\right), \tag{5}$$

with G^μ_ν the Einstein tensor and $\kappa^2 \equiv 8\pi G = 8\pi/M_P^2$. From these formulae (4) and (5) we obtain the background equation of motion for the scalar fields $\boldsymbol{\phi}$,

$$\mathcal{D}\boldsymbol{\phi}^{;} + 3H_a\boldsymbol{\phi}^{;} + b^2\,\mathbf{G}^{-1}\boldsymbol{\nabla}^T V = 0, \tag{6}$$

and the Friedmann equations

$$H_a^2 = \frac{1}{3}\kappa^2\left(\frac{1}{2}|\boldsymbol{\phi}^{;}|^2 + b^2 V\right), \qquad\qquad \mathcal{D}H_a = -\frac{1}{2}\kappa^2|\boldsymbol{\phi}^{;}|^2. \tag{7}$$

Here we have introduced the "slow-roll derivative" $\mathcal{D}$ which is defined as follows: on any quantity A that does not have any b dependence, $\mathcal{D}(b^n A) = (\mathcal{D}_\tau - nH_b)(b^n A)$. In particular this means that $\mathcal{D}\dot{\boldsymbol{\phi}} = (\mathcal{D}_\tau - H_b)\dot{\boldsymbol{\phi}}$, $\mathcal{D}^2\dot{\boldsymbol{\phi}} = (\mathcal{D}_\tau - 2H_b)(\mathcal{D}_\tau - H_b)\dot{\boldsymbol{\phi}}$, $\mathcal{D}H_a = (\partial_\tau - H_b)H_a$, etc. Notice that the slow-roll derivative equals the comoving time derivative $\mathcal{D}_t$ if comoving time is used ($b = 1$), while with conformal time it reads $\mathcal{D} = \mathcal{D}_\eta - n\mathcal{H}$. The slow-roll derivative is a necessary ingredient to be able to write these equations in terms of a general time variable. It has the important property that when it is applied to quantities like field velocities or Hubble parameters, only terms of one order higher in the slow-roll approximation are obtained (hence its name), as we show in the next section.

We finish this section by introducing a prefered basis $\{\boldsymbol{e}_n\}$ on the field manifold that is induced by the dynamics of the system. This basis was already introduced in our previous paper [37], although not in the context of a general time variable. The treatment with the angle of [18] is a special case of this basis in the limit of only two fields. The first unit vector $\boldsymbol{e}_1$ is given by the direction of the field velocity $\dot{\boldsymbol{\phi}}$. The second unit vector $\boldsymbol{e}_2$ points in the direction of that part of the field acceleration $\mathcal{D}\dot{\boldsymbol{\phi}}$ that is perpendicular to the first unit vector $\boldsymbol{e}_1$. This Gram-Schmidt orthogonalization process can be extended to any n: the unit vector $\boldsymbol{e}_n$ points in the direction of $\boldsymbol{\phi}^{(n)} \equiv \mathcal{D}^{(n-1)}\dot{\boldsymbol{\phi}}$ that is perpendicular to the first $n-1$ unit vectors $\boldsymbol{e}_1, \ldots, \boldsymbol{e}_{n-1}$. Using the projection operators $\mathbf{P}_n$, which project on the $\boldsymbol{e}_n$, and $\mathbf{P}_n^\perp$, which project on the subspace that is perpendicular to $\boldsymbol{e}_1, \ldots, \boldsymbol{e}_n$, the definitions of the unit vectors are given by

$$\boldsymbol{e}_n = \frac{\mathbf{P}_{n-1}^\perp \boldsymbol{\phi}^{(n)}}{|\mathbf{P}_{n-1}^\perp \boldsymbol{\phi}^{(n)}|}, \qquad \mathbf{P}_n = \boldsymbol{e}_n \boldsymbol{e}_n^\dagger, \qquad \mathbf{P}_n^\perp = \mathbb{1} - \sum_{q=1}^{n} \mathbf{P}_q, \tag{8}$$

for all $n = 1, 2, \ldots$ and with the definition $\mathbf{P}_0^\perp \equiv \mathbb{1}$. Notice that the unit vectors $\boldsymbol{e}_n$ will in general depend on time. However, because the slow-roll derivative $\mathcal{D}$ was used in the definition of this basis, the definition does not depend on a specific choice of time variable. By construction the vector $\boldsymbol{\phi}^{(n)}$ can be expanded in these unit vectors as

$$\boldsymbol{\phi}^{(n)} = (\mathbf{P}_1 + \ldots + \mathbf{P}_n)\, \boldsymbol{\phi}^{(n)} = \sum_{p=1}^{n} \phi^{(n)}_{\ p} \boldsymbol{e}_p, \qquad \phi^{(n)}_{\ p} = \boldsymbol{e}_p \cdot \boldsymbol{\phi}^{(n)}. \tag{9}$$

In particular, we have that $\phi^{(n)}_{\ n} = \boldsymbol{e}_n \cdot \boldsymbol{\phi}^{(n)} = |\mathbf{P}_{n-1}^\perp \boldsymbol{\phi}^{(n)}|$. As the projection operators $\mathbf{P}_1$ and $\mathbf{P}_1^\perp$ turn out to be the most important in our discussions, we introduce the short-hand notation $\mathbf{P}^\parallel = \mathbf{P}_1$ and $\mathbf{P}^\perp = \mathbf{P}_1^\perp = \mathbb{1} - \mathbf{P}^\parallel$. In terms of these two operators we can write a general vector and matrix as $\boldsymbol{A} = \boldsymbol{A}^\parallel + \boldsymbol{A}^\perp$ and $\mathbf{M} = \mathbf{M}^{\parallel\parallel} + \mathbf{M}^{\parallel\perp} + \mathbf{M}^{\perp\parallel} + \mathbf{M}^{\perp\perp}$, with $\boldsymbol{A}^\parallel \equiv \mathbf{P}^\parallel \boldsymbol{A}$ and $\mathbf{M}^{\parallel\parallel} \equiv \mathbf{P}^\parallel \mathbf{M} \mathbf{P}^\parallel$, etc.

2.2. Slow roll

Slow-roll inflation is driven by a scalar field potential that is almost flat and therefore acts as an effective cosmological constant. In the case of a single scalar field, the notion of slow roll is well-established (see e.g. [7, 19, 38]). This concept can be generalized to multiple scalar fields in a geometrical way using the unit vectors introduced in the previous section. The system consisting of (6) and (7) is said to be in the slow-roll regime if the comoving time derivatives satisfy $|\mathcal{D}_t\dot{\boldsymbol{\phi}}| \ll |3H\dot{\boldsymbol{\phi}}|$ and $\frac{1}{2}|\dot{\boldsymbol{\phi}}|^2 \ll V$. A more precise definition not depending on the use of comoving time is given below (14).

We introduce the following functions for an arbitrary time variable τ (see also [37]):

$$\tilde{\epsilon}(\boldsymbol{\phi}) \equiv -\frac{\mathcal{D}H_a}{H_a^2}, \qquad \tilde{\boldsymbol{\eta}}^{(n)}(\boldsymbol{\phi}) \equiv \frac{\mathcal{D}^{n-1}\boldsymbol{\phi}^{;}}{(H_a)^{n-1}|\boldsymbol{\phi}^{;}|}. \tag{10}$$

We often use the short-hand notation $\tilde{\boldsymbol{\eta}} = \tilde{\boldsymbol{\eta}}^{(2)}$ and $\tilde{\boldsymbol{\xi}} = \tilde{\boldsymbol{\eta}}^{(3)}$. Both these vectors can be decomposed in components parallel $(\tilde{\eta}^{\|}, \tilde{\xi}^{\|})$ and perpendicular $(\tilde{\eta}^{\perp})$ to the field velocity $\boldsymbol{\phi}^{;}$:

$$\tilde{\eta}^{\|} = \boldsymbol{e}_1 \cdot \tilde{\boldsymbol{\eta}} = \frac{\mathcal{D}\boldsymbol{\phi}^{;} \cdot \boldsymbol{\phi}^{;}}{H_a|\boldsymbol{\phi}^{;}|^2}, \qquad \tilde{\eta}^{\perp} = \boldsymbol{e}_2 \cdot \tilde{\boldsymbol{\eta}} = \frac{|(\mathcal{D}\boldsymbol{\phi}^{;})^{\perp}|}{H_a|\boldsymbol{\phi}^{;}|}, \qquad \tilde{\xi}^{\|} = \boldsymbol{e}_1 \cdot \tilde{\boldsymbol{\xi}} = \frac{\mathcal{D}^2\boldsymbol{\phi}^{;} \cdot \boldsymbol{\phi}^{;}}{H_a^2|\boldsymbol{\phi}^{;}|^2}. \tag{11}$$

(Even though $\tilde{\boldsymbol{\xi}}$ in general has two directions perpendicular to $\boldsymbol{e}_1$ we only give $\xi^{\|}$ here since it is the only one that turns out to be relevant in the remainder of this work.) The derivatives of the slow-roll functions can be computed from their definitions and are given by:

$$\tilde{\epsilon}^{;} = 2H_a\tilde{\epsilon}(\tilde{\epsilon} + \tilde{\eta}^{\|}), \quad (\tilde{\eta}^{\|})^{;} = H_a[\tilde{\xi}^{\|} + (\tilde{\eta}^{\perp})^2 + \tilde{\epsilon}\tilde{\eta}^{\|} - (\tilde{\eta}^{\|})^2], \ \mathcal{D}\tilde{\boldsymbol{\eta}} = H_a[\tilde{\boldsymbol{\xi}} + (\tilde{\epsilon} - \tilde{\eta}^{\|})\tilde{\boldsymbol{\eta}}]. \tag{12}$$

In terms of the functions $\tilde{\epsilon}, \tilde{\boldsymbol{\eta}}$ the Friedmann equation (7) and the background field equation (6) read

$$H_a = \frac{\kappa}{\sqrt{3}}\, b\sqrt{V}\left(1 - \frac{1}{3}\tilde{\epsilon}\right)^{-1/2}, \tag{13}$$

$$\boldsymbol{\phi}^{;} + \frac{2}{\sqrt{3}}\frac{1}{\kappa}\, b\,\mathbf{G}^{-1}\boldsymbol{\nabla}^T\sqrt{V} = -\sqrt{\frac{2}{3}}\, b\sqrt{V}\,\frac{\sqrt{\tilde{\epsilon}}}{1 - \frac{1}{3}\tilde{\epsilon}}\left(\frac{1}{3}\tilde{\boldsymbol{\eta}} + \frac{\frac{1}{3}\tilde{\epsilon}\,\boldsymbol{e}_1}{1 + \sqrt{1 - \frac{1}{3}\tilde{\epsilon}}}\right). \tag{14}$$

(Notice that for a positive potential V the function $\tilde{\epsilon} < 3$, as can be seen from its definition.) We can now define precisely what is meant by slow roll as these two background equations are still exact. Slow roll is valid if $\tilde{\epsilon}$, $\sqrt{\tilde{\epsilon}}\,\tilde{\eta}^{\|}$ and $\sqrt{\tilde{\epsilon}}\,\tilde{\eta}^{\perp}$ are (much) smaller than unity. For this reason $\tilde{\epsilon}$, $\tilde{\eta}^{\|}$ and $\tilde{\eta}^{\perp}$ are called slow-roll functions. The function $\tilde{\boldsymbol{\xi}}$ is called a second order slow-roll function because it involves two slow-roll derivatives, and it is assumed to be of an order comparable to $\tilde{\epsilon}^2$, $\tilde{\epsilon}\tilde{\eta}^{\|}$, etc. If slow roll is valid, we can use expansions in powers of these slow-roll functions to estimate the relevance of various terms in a given expression. For example, to first order the Friedmann equation (13) is approximated by replacing $(1 - \tilde{\epsilon}/3)^{-1/2}$ by $(1 + \tilde{\epsilon}/6)$. The background field equation up to and including first order is given by (14) with the right-hand side set to zero, as all those terms are order 3/2 or higher. This last fact is the motivation for defining results accurate to first order to have no corrections larger than order $\tilde{\epsilon}^{3/2}$ when discussing the perturbations in section 3.3.

At the level of the solutions of these equations we make the following definition. An approximate solution of an equation of motion is said to be accurate to first order in slow roll, if the relative difference between this solution and the exact one is of a smaller numerical order than the slow-roll functions. This relative error depends in general on the size of the integration interval. Let us explain this with the following example. From (12) we see that the time derivatives of the slow-roll functions are second order quantities. Hence we can make the assumption that to first order the slow-roll functions are constant. Switching to the number of e-folds N, which is related as $\mathrm{d}N = H_a \mathrm{d}\tau$ to the time variable τ, we can integrate (12) to find the variation of $\tilde{\epsilon}$ over an interval $[N_1, N_2]$:

$$\Delta\tilde{\epsilon} = \int_{N_1}^{N_2} \mathrm{d}N \, 2\tilde{\epsilon}(\tilde{\epsilon} + \tilde{\eta}^{\parallel}) = 2\tilde{\epsilon}_0(\tilde{\epsilon}_0 + \tilde{\eta}_0^{\parallel})(N_2 - N_1). \tag{15}$$

Here the subscript $_0$ denotes some reference time in this interval where the slow-roll functions are evaluated. Hence we see that if the interval $(N_2 - N_1)$ becomes larger than $1/(2(\tilde{\epsilon}_0 + \tilde{\eta}_0^{\parallel}))$, $\Delta\tilde{\epsilon}$ becomes larger than $\tilde{\epsilon}_0$ and the assumption of taking $\tilde{\epsilon}$ constant over this interval to first order is certainly not valid anymore. (An example of the real behaviour of the slow-roll functions can be found in figure 1b) in section 4.2.) In the literature these effects are usually ignored and the solution of an equation of motion valid to first order is (implicitly) assumed to be accurate to first order as well. However, with that assumption the numerical error between slow-roll and exact solution can become very large depending on the size of the interval of integration, which is the reason for our revised definition.

The slow-roll functions (10) are all defined as functions of covariant derivatives of the velocity $\boldsymbol{\phi}^{;}$ and the Hubble parameter H_a. If the zeroth order slow-roll approximation works well, that is if the right-hand side of (14) can be neglected, as well as the $\tilde{\epsilon}$ in (13), then we can use these two equations to eliminate $\boldsymbol{\phi}^{;}$ and H_a in favour of the potential V. This is the way the original single-field slow-roll parameters were defined. However, that original definition had the disadvantage that the slow-roll conditions became consistency checks. While we can expand the exact equations in powers of the slow-roll functions, that is impossible by construction with the original slow-roll parameters in terms of the potential.‡ In order to avoid confusion we compare the slow-roll functions we defined in (10) with the ones originally used in the single-field case, ϵ and η:

$$\epsilon = \frac{1}{2\kappa^2} \frac{V_{,\phi}^2}{V^2} = \tilde{\epsilon}, \qquad\qquad \eta = \frac{1}{\kappa^2} \frac{V_{,\phi\phi}}{V} = -\tilde{\eta}^{\parallel} + \tilde{\epsilon}, \tag{16}$$

where the last equalities in both equations are only valid to lowest order in the slow-roll approximation.

For later use we introduce the matrix Z by

$$(Z)_{mn} = -(Z^T)_{mn} = \frac{1}{H_a} \boldsymbol{e}_m^{\dagger} \mathcal{D} \boldsymbol{e}_n, \tag{17}$$

which shows a nice interplay between the unit vectors and the notion of slow roll. The anti-symmetry of Z follows because $(\boldsymbol{e}_m^{\dagger} \boldsymbol{e}_n)^{;} = 0$. To determine its components we observe that

$$\mathcal{D}\boldsymbol{e}_{n+1} \cdot \boldsymbol{e}_{n-p} + \boldsymbol{e}_{n+1} \cdot \mathcal{D}\boldsymbol{e}_{n-p} = 0, \qquad \mathcal{D}\boldsymbol{e}_{n+1} \cdot \tilde{\boldsymbol{\eta}}^{(n)} + H_a \, \boldsymbol{e}_{n+1} \cdot \tilde{\boldsymbol{\eta}}^{(n+1)} = 0, \tag{18}$$

---

‡ In the context of single-field inflation this was discussed in detail in [38].

because $\boldsymbol{e}_{n+1}$ is perpendicular to $\boldsymbol{e}_{n-p}$ with $0 \leq p < n$ and to $\tilde{\boldsymbol{\eta}}^{(n)}$. From the construction of $\boldsymbol{\phi}^{(n)}$ in (9) we see that $\mathcal{D}\boldsymbol{e}_n$ can never get a component in a direction higher than $\boldsymbol{e}_{n+1}$. Hence we deduce from the first equation in (18) that for $p \geq 1$, $\mathcal{D}\boldsymbol{e}_{n+1}$ and $\boldsymbol{e}_{n-p}$ are perpendicular. Using this we see that of the first term of the second equation only the $\boldsymbol{e}_n$ direction is relevant, so that the only non-zero components of Z read

$$Z_{n\,n+1} = -Z_{n+1\,n} = -\frac{\boldsymbol{e}_{n+1} \cdot \tilde{\boldsymbol{\eta}}^{(n+1)}}{\boldsymbol{e}_n \cdot \tilde{\boldsymbol{\eta}}^{(n)}}, \tag{19}$$

which is first order in slow roll.

3. Perturbations in multiple-field inflation

3.1. Equations of motion for the perturbations

This section describes the coupled system of gravity, encoded by the metric $g_{\mu\nu}$, and multiple scalar field perturbations $\boldsymbol{\delta\phi}$ during inflation. We separate both the scalar fields and the metric into a homogeneous background part and an inhomogeneous perturbation, which is assumed to be small. Since the observed fluctuations in the CMBR are tiny, this assumption is well-motivated. Consequently one can linearize all equations with respect to the perturbations. We define

$$\begin{aligned}&\boldsymbol{\phi}^{\text{full}}(\eta, \boldsymbol{x}) = \boldsymbol{\phi}(\eta) + \boldsymbol{\delta\phi}(\eta, \boldsymbol{x}),\\ &g^{\text{full}}_{\mu\nu}(\eta, \boldsymbol{x}) = g_{\mu\nu}(\eta) + \delta g_{\mu\nu}(\eta, \boldsymbol{x}) = a^2(\eta)\begin{pmatrix}-1 & 0\\ 0 & \delta_{ij}\end{pmatrix} - 2a^2(\eta)\Phi(\eta, \boldsymbol{x})\begin{pmatrix}1 & 0\\ 0 & \delta_{ij}\end{pmatrix}.\end{aligned} \tag{20}$$

As is discussed in [6], this metric is obtained by applying the so-called longitudinal gauge to the flat Robertson-Walker metric in the case when only scalar metric perturbations and a scalar matter theory are considered. In this gauge all formulae look the same as when the gauge-invariant approach [39, 6] is used. The gravitational (Newtonian) potential $\Phi(\eta, \boldsymbol{x})$ describes the scalar metric perturbations.

The equation of motion for the perturbations of the metric is obtained by linearizing and combining the (00) and (ii) components of the Einstein equations (5):

$$\Phi'' + 6\mathcal{H}\Phi' + 2(\mathcal{H}' + 2\mathcal{H}^2)\Phi - \Delta\Phi = -\kappa^2 a^2(\boldsymbol{\nabla} V\, \boldsymbol{\delta\phi}), \tag{21}$$

where the spatial Laplacean is given by $\Delta = \sum_i \partial_i^2$, while the integrated $(0i)$ component of the Einstein equations leads to the constraint equation

$$\Phi' + \mathcal{H}\Phi = \frac{1}{2}\kappa^2 \boldsymbol{\phi}' \cdot \boldsymbol{\delta\phi} = \frac{1}{2}\kappa^2 |\boldsymbol{\phi}'| \delta\phi^{\parallel}. \tag{22}$$

Here we have decomposed $\boldsymbol{\delta\phi} = \delta\phi^{\parallel}\, \boldsymbol{e}_1 + \boldsymbol{\delta\phi}^{\perp}$. (A similar decomposition in the case of two-field inflation was also discussed in [18].) In addition we have the equation of motion for the scalar field perturbations,

$$\Big(\mathcal{D}_\eta^2 + 2\mathcal{H}\mathcal{D}_\eta - \Delta + a^2 \tilde{\mathbf{M}}^2(\boldsymbol{\phi})\Big)\, \boldsymbol{\delta\phi} = 4\Phi'\boldsymbol{\phi}' - 2a^2\Phi\, \mathbf{G}^{-1}\boldsymbol{\nabla}^T V, \tag{23}$$

where we have introduced the (effective) mass-matrices

$$\tilde{\mathbf{M}}^2 \equiv \mathbf{M}^2 - \mathbf{R}(\dot{\boldsymbol{\phi}}, \dot{\boldsymbol{\phi}}), \qquad \mathbf{M}^2 \equiv \mathbf{G}^{-1}\boldsymbol{\nabla}^T\boldsymbol{\nabla} V, \tag{24}$$

with $\mathbf{R}$ the field curvature as defined in the appendix. This system of perturbation equations must be solved in the background determined by the scalar fields (6) and the Friedmann equations (7). Using the integrated $(0i)$ component of the Einstein equations (22) together with the background equation of motion for the scalar fields (6), the right-hand side of equation (21) for Φ can be rewritten as

$$-\kappa^2 a^2 (\boldsymbol{\nabla} V\, \boldsymbol{\delta\phi}) = 2(\Phi' + \mathcal{H}\Phi)\left(\frac{1}{|\boldsymbol{\phi}'|}(\mathcal{D}_\eta \boldsymbol{\phi}') \cdot \boldsymbol{e}_1 + 2\mathcal{H}\right) + \kappa^2 (\mathcal{D}_\eta \boldsymbol{\phi}') \cdot \boldsymbol{\delta\phi}^\perp, \tag{25}$$

where we used the definition of the projection operators. Inserting this expression in (21) and realizing that $|\boldsymbol{\phi}'|' \, |\boldsymbol{\phi}'| = (\mathcal{D}_\eta \boldsymbol{\phi}') \cdot \boldsymbol{\phi}'$, we get

$$\Phi'' + 2\left(\mathcal{H} - \frac{|\boldsymbol{\phi}'|'}{|\boldsymbol{\phi}'|}\right)\Phi' + 2\left(\mathcal{H}' - \mathcal{H}\frac{|\boldsymbol{\phi}'|'}{|\boldsymbol{\phi}'|}\right)\Phi - \Delta\Phi = \kappa^2 (\mathcal{D}_\eta \boldsymbol{\phi}') \cdot \boldsymbol{\delta\phi}^\perp. \tag{26}$$

In the single-field case the right-hand side is zero because $\boldsymbol{\delta\phi}^\perp$ then vanishes by construction.

The system of perturbations (26), (22) and (23) is quite complicated. To make the physical content more transparent, we introduce new variables u and $\boldsymbol{q}$ (linearly related to Φ and $\boldsymbol{\delta\phi}$, respectively),

$$u \equiv \frac{a}{\kappa^2 |\boldsymbol{\phi}'|}\,\Phi = \frac{\Phi}{\kappa\sqrt{2}\, H\sqrt{\tilde{\epsilon}}}, \qquad \boldsymbol{q} \equiv a\left(\boldsymbol{\delta\phi} + \frac{\Phi}{\mathcal{H}}\,\boldsymbol{\phi}'\right), \tag{27}$$

which satisfy the following two requirements:

(i) The equations of motion for both u and $\boldsymbol{q}$ do not contain first order conformal time derivatives;

(ii) The equation of motion for $\boldsymbol{q}$ is homogeneous and $\boldsymbol{q}$ is gauge invariant.

The first requirement makes a direct comparison between the size of the Fourier mode $\boldsymbol{k}^2 = k^2$ and other physical background quantities in the equation of motion possible. In section 3.3 we make use of this to distinguish between different regions for the behaviour of the solutions. The other requirement ensures that we can naively quantize $\boldsymbol{q}$ using the Lagrangean corresponding to the equation of motion for $\boldsymbol{q}$ in section 3.2. As $\boldsymbol{q}$ is gauge invariant and linearly related to $\boldsymbol{\delta\phi}$, apart from the shift proportional to $\Phi\boldsymbol{\phi}'$, no non-physical degrees of freedom are quantized. The single-field version of $\boldsymbol{q}$, including its equation of motion and quantization, was first introduced by Sasaki and Mukhanov [40, 41], which is why variables of this type are sometimes refered to as Sasaki-Mukhanov variables. The variable u was first introduced by Mukhanov [42], see also [6].

To derive the equation of motion for $\boldsymbol{q}$ we need an auxiliary result. By differentiating the background field equation in terms of conformal time, $\mathcal{D}_\eta \boldsymbol{\phi}' + 2\mathcal{H}\boldsymbol{\phi}' + a^2 \mathbf{G}^{-1}\boldsymbol{\nabla}^T V = 0$, once more we obtain

$$\mathcal{D}_\eta^2 \boldsymbol{\phi}' + 2(\mathcal{H}' - 2\mathcal{H}^2)\boldsymbol{\phi}' + a^2 \tilde{\mathbf{M}}^2 \boldsymbol{\phi}' = 0, \tag{28}$$

where we used that $\mathcal{D}_\eta(\mathbf{G}^{-1}\boldsymbol{\nabla}^T V) = \mathbf{M}^2 \boldsymbol{\phi}' = \tilde{\mathbf{M}}^2 \boldsymbol{\phi}'$ (because of the anti-symmetry properties of the curvature tensor $\mathbf{R}(\dot{\boldsymbol{\phi}}, \dot{\boldsymbol{\phi}})\boldsymbol{\phi}' = 0$). The equation for $\boldsymbol{q}$ is then obtained from the equation of motion (23) for $\boldsymbol{\delta\phi}$ and (21), (22) for Φ, using the projectors (8) and slow-roll functions (10). Combining this with the derivatives of $\mathcal{H}$ from (7) and of the slow-roll functions (12), we finally obtain the homogeneous equation for the spatial Fourier mode $\boldsymbol{k}$ of $\boldsymbol{q}$:

$$\mathcal{D}_\eta^2 \boldsymbol{q}_{\boldsymbol{k}} + (k^2 + \mathcal{H}^2 \boldsymbol{\Omega}) \boldsymbol{q}_{\boldsymbol{k}} = 0, \qquad L = \frac{1}{2} \mathcal{D}_\eta \boldsymbol{q}_{\boldsymbol{k}}^\dagger \mathcal{D}_\eta \boldsymbol{q}_{\boldsymbol{k}} - \frac{1}{2} \boldsymbol{q}_{\boldsymbol{k}}^\dagger (k^2 + \mathcal{H}^2 \boldsymbol{\Omega}) \boldsymbol{q}_{\boldsymbol{k}}. \tag{29}$$

Here L is the associated Lagrangean and

$$\boldsymbol{\Omega} \equiv \frac{1}{H^2} \tilde{\mathbf{M}}^2 - (2 - \tilde{\epsilon}) \mathbb{1} - 2\tilde{\epsilon} \Big((3 + \tilde{\epsilon}) \mathbf{P}^\parallel + \boldsymbol{e}_1 \tilde{\boldsymbol{\eta}}^\dagger + \tilde{\boldsymbol{\eta}} \boldsymbol{e}_1^\dagger \Big). \tag{30}$$

The $(n1)$ components of $\boldsymbol{\Omega}$ can be expressed completely in terms of slow-roll functions using

$$\frac{1}{H^2} \tilde{\mathbf{M}}^2 \boldsymbol{e}_1 = \frac{1}{H^2} \mathbf{M}^2 \boldsymbol{e}_1 = 3\, \tilde{\epsilon}\, \boldsymbol{e}_1 - 3\, \tilde{\boldsymbol{\eta}} - \tilde{\boldsymbol{\xi}}. \tag{31}$$

The other components can in general not be expressed in terms of the slow-roll functions introduced in the previous subsection.

To derive the equation of motion for u it is convenient to introduce the quantity θ,

$$\theta \equiv \frac{\mathcal{H}}{a |\boldsymbol{\phi}'|} = \frac{\kappa}{\sqrt{2}} \frac{1}{a \sqrt{\tilde{\epsilon}}} \tag{32}$$

$$\Rightarrow \qquad \frac{\theta'}{\theta} = -\mathcal{H} \left(1 + \tilde{\epsilon} + \tilde{\eta}^\parallel \right), \qquad \frac{\theta''}{\theta} = \mathcal{H}^2 \left(2\tilde{\epsilon} + \tilde{\eta}^\parallel + 2 (\tilde{\eta}^\parallel)^2 - (\tilde{\eta}^\perp)^2 - \tilde{\xi}^\parallel \right),$$

where we also gave the resulting expressions for its derivatives, and observe that the following relations hold for the slow-roll functions:

$$\mathcal{H}' = \mathcal{H}^2 (1 - \tilde{\epsilon}), \qquad \frac{|\boldsymbol{\phi}'|'}{|\boldsymbol{\phi}'|} = \mathcal{H} (1 + \tilde{\eta}^\parallel), \qquad (\mathcal{D}_\eta \boldsymbol{\phi}')^\perp = \frac{\sqrt{2}}{\kappa} \mathcal{H}^2 \sqrt{\tilde{\epsilon}}\, \tilde{\boldsymbol{\eta}}^\perp. \tag{33}$$

By substituting the definitions of u and $\boldsymbol{q}$ in (26), where we first rewrite the relation between Φ and u as $\Phi = \kappa \sqrt{2} \mathcal{H} \sqrt{\tilde{\epsilon}}\, u / a$, and using the above expressions and the derivatives of the slow-roll functions given in (12), we obtain

$$u_{\boldsymbol{k}}'' + \left(k^2 - \frac{\theta''}{\theta} \right) u_{\boldsymbol{k}} = \mathcal{H} \tilde{\eta}^\perp \boldsymbol{e}_2 \cdot \boldsymbol{q}_{\boldsymbol{k}}. \tag{34}$$

Notice that all these equations are still exact, no slow-roll approximations have been made. From this one can draw the conclusion that at the level of the equations the redefined gravitational potential u decouples from the perpendicular components of the field perturbation $\boldsymbol{q}^\perp$ to leading order, but at first order mixing between these perturbations appears.

The equations of motion (29) and (34) show that the different spatial Fourier modes of both $\boldsymbol{q}$ and u decouple. From now on we only consider one generic mode $\boldsymbol{k}$, so that we can drop the subscripts $_{\boldsymbol{k}}$. Rewriting equation (22) in terms of the components $q_n \equiv \boldsymbol{e}_n \cdot \boldsymbol{q}$ of $\boldsymbol{q}$ and differentiating it once gives

$$u' - \frac{\theta'}{\theta} u = \frac{1}{2} q_1 \qquad \Rightarrow \qquad u'' - \frac{\theta''}{\theta} u = \frac{1}{2} \Big(q_1' + \frac{\theta'}{\theta} q_1 \Big), \tag{35}$$

where θ and its derivatives are given in (32). This equation for u'' can be combined with the equation of motion (34) for u to give

$$k^2 u = \mathcal{H} \tilde{\eta}^\perp q_2 - \frac{1}{2} \Big(q_1' + \frac{\theta'}{\theta} q_1 \Big). \tag{36}$$

After $\boldsymbol{q}$ has been quantized, this expression can be used to relate it to u. (Although this relation could in principle be used to compute u at the end of inflation, its numerical implementation can be rather awkward because of cancellation of large numbers. In numerical situations it turns out to be more convenient to determine u from its own equation of motion and only use (36) to find the correct quantization and initial conditions.)

3.2. Quantization of the perturbations

We start with the Lagrangean (29) in terms of the basis $\{\boldsymbol{e}_n\}$:

$$L = \frac{1}{2}(q' + \mathcal{H}Zq)^T(q' + \mathcal{H}Zq) - \frac{1}{2}q^T(k^2 + \mathcal{H}^2\Omega)q, \tag{37}$$

where we employ the notation $(\Omega)_{mn} = \boldsymbol{e}_m^\dagger \boldsymbol{\Omega} \boldsymbol{e}_n$ and the matrix Z is given in (17). Notice that this Lagrangean has the standard canonical normalization of $\frac{1}{2}(q')^T q'$, independent of the field metric $\mathbf{G}$, as can be derived from the original Lagrangean (2). We maintain the vectorial structure of this multiple-field system and repress the indices n, m as much as possible, which means for example that the non-bold q in this equation is a vector (in the basis $\{\boldsymbol{e}_n\}$). From the canonical momenta $\pi = \partial L/\partial q'^T$ we find the Hamiltonian $H = \pi^T q' - L$ and the Hamilton equations:

$$\begin{aligned} H &= \frac{1}{2}(\pi - \mathcal{H}Zq)^T(\pi - \mathcal{H}Zq) + \frac{1}{2}q^T\Big(k^2 + \mathcal{H}^2(\Omega + Z^2)\Big)q; \\ q' &= \frac{\partial H}{\partial \pi^T} = \pi - \mathcal{H}Zq, \qquad \pi' = -\frac{\partial H}{\partial q^T} = -(k^2 + \mathcal{H}^2\Omega)q - \mathcal{H}Z\pi. \end{aligned} \tag{38}$$

In order to avoid writing indices when considering commutation relations we use vectors α, β with components α_m, β_m in the $\boldsymbol{e}_m$ basis that are independent of q and π. The canonical commutation relations can then be represented as

$$[\alpha^T\hat{q}, \beta^T\hat{q}] = [\alpha^T\hat{\pi}, \beta^T\hat{\pi}] = 0, \qquad [\alpha^T\hat{q}, \beta^T\hat{\pi}] = i\alpha^T\beta. \tag{39}$$

Using the Hamilton equations it can be checked that this quantization procedure is indeed time independent. Let Q and Π be complex matrix valued solutions of the Hamilton equations, such that $q = Qa_0^* + \text{c.c.}$, $\pi = \Pi a_0^* + \text{c.c.}$ is a solution of (38) for any constant complex vector a_0. Here c.c. denotes the complex conjugate. The Hamilton equations for Q and Π can be combined to give a second order differential equation for Q. To remove the first order time derivative from this equation, we define $Q(\eta) = R(\eta)\tilde{Q}(\eta)$ with R chosen such that the matrix functions R and $\tilde{Q}$ satisfy

$$R' + \mathcal{H}ZR = 0, \qquad \tilde{Q}'' + (k^2 + \mathcal{H}^2\tilde{\Omega})\tilde{Q} = 0, \qquad \text{with} \quad \tilde{\Omega} = R^{-1}\Omega R. \tag{40}$$

The matrix Π is then given by $\Pi = Q' + \mathcal{H}ZQ = R\tilde{Q}'$. We take $R(\eta_i) = \mathbb{1}$ as initial condition, since the initial condition of Q can be absorbed in that of $\tilde{Q}$. The equation of motion for R implies that $R^T R$ and $\ln\det R$ are constant because Z is anti-symmetric and consequently traceless. Taking into account its initial condition, it then follows that R represents a rotation.

It now follows that $\hat{q}$ and $\hat{\pi}$ can be expanded in terms of constant creation ($\hat{a}^\dagger$) and annihilation ($\hat{a}$) operator vectors:

$$\hat{q} = Q\hat{a}^\dagger + Q^*\hat{a} = R\tilde{Q}\hat{a}^\dagger + R\tilde{Q}^*\hat{a}, \qquad \hat{\pi} = \Pi\hat{a}^\dagger + \Pi^*\hat{a}. \tag{41}$$

The creation and annihilation operators satisfy

$$[\alpha^T\hat{a}, \beta^T\hat{a}] = [\alpha^T\hat{a}^\dagger, \beta^T\hat{a}^\dagger] = 0, \qquad [\alpha^T\hat{a}, \beta^T\hat{a}^\dagger] = \alpha^T\beta. \tag{42}$$

This is consistent with the commutation relations for q and π given above, provided that the matrix functions Q and Π satisfy

$$Q^*\,Q^T - Q\,Q^{*T} = \Pi^*\,\Pi^T - \Pi\,\Pi^{*T} = 0, \qquad Q^*\,\Pi^T - Q\,\Pi^{*T} = i\mathbb{1}. \tag{43}$$

These relations hold for all time, as can be checked explicitly by using the equations of motion for Q and Π to show that they are time independent, provided that they hold at some given time.

We assume that the initial state is the vacuum $|0\rangle$ defined by $\hat{a}|0\rangle = 0$ and that there is no initial particle production. This implies that the Hamiltonian initially does not contain any terms with $\hat{a}\hat{a}$ and $\hat{a}^\dagger\hat{a}^\dagger$, which leads to the condition

$$(\Pi - \mathcal{H}ZQ)^T(\Pi - \mathcal{H}ZQ) + Q^T\Big(k^2 + \mathcal{H}^2(\Omega - Z^TZ)\Big)Q = 0. \tag{44}$$

The solution of the equations (43) and (44) can be parametrized by a unitary matrix U at the beginning of inflation, when the limit that k^2 is much bigger than any other scale is applicable:

$$Q_i = \frac{1}{\sqrt{2k}}U, \qquad \Pi_i = \frac{i\sqrt{k}}{\sqrt{2}}U. \tag{45}$$

We denote expectation values with respect to the vacuum state $|0\rangle$ by $\langle\ldots\rangle$. Let α, β be two vectors. Then for the expectation value of $(\alpha^TQU\hat{a}^\dagger + \alpha^{*T}Q^*U^*\hat{a})^2$, with U a unitary matrix, we obtain

$$\langle(\alpha^TQU\hat{a}^\dagger + \alpha^{*T}Q^*U^*\hat{a})^2\rangle = \alpha^{*T}Q^*U^*U^TQ^T\alpha = \alpha^{*T}Q^*Q^T\alpha. \tag{46}$$

So a unitary matrix in front of the $\hat{a}^\dagger$ will drop out in the computation of this correlator. This is even true if another state than the vacuum is used to compute the correlator. In particular this means that the correlator of the gravitational potential will not depend on the unitary matrix U in (45). To draw this conclusion we use relation (36) between u and $\boldsymbol{q}$ and the fact that Q satisfies a linear homogeneous equation of motion. We also see that as long as Q is simply oscillating and hence itself unitary (apart from a normalization factor), its evolution will be irrelevant for the computation of the correlator.

We finish this section with some brief remarks on the assumption of taking the vacuum state to compute the correlator. The vacuum state $|0\rangle$ at the beginning of inflation seems a reasonable assumption for the calculation of the density perturbations that we can observe in the CMBR today. Even though perturbations in the CMBR have long wavelengths now, they had very short wavelengths before they crossed the Hubble radius during inflation. Therefore, their scale k at the beginning of inflation at t_i is much larger than the Planck scale. It seems a reasonable assumption that modes with momenta very much larger than the Planck scale are not excited at t_i, so that for these modes the vacuum state is a good assumption.§ This assumption can be tested by taking other states than the vacuum state. For instance one can try a thermal state with a temperature of the Planck scale. Typically one finds that if there were a few e-folds of inflation before the now observable scales crossed the Hubble scale, corrections are negligible. For a more detailed discussion on observable effects of non-vacuum initial states we refer to [45, 46].

3.3. Solutions of the perturbation equations to first order: setup

To derive analytical expressions for the gravitational potential and field perturbations valid to first order in slow roll, we have to determine the evolution of the modified Newtonian potential u and quantized variables $\boldsymbol{q}$, described by the equations (34), (29) and (35), analytically and accurately up to first order during inflation. In this section we explain the physical ideas that go into that computation. Section 3.4 introduces the concept of slow roll on the perturbations, which is useful in part of the calculations that are presented in section 3.5. The treatment here has been partly inspired by the discussion of the transition region in [9].

Since $\mathcal{H}$ grows rapidly, while k is constant for a given mode, the solutions of (34) and (29) change dramatically around the time $\eta_\mathcal{H}$ when a scale crosses the Hubble scale. This time is defined by the relation

$$\mathcal{H}(\eta_\mathcal{H}) = k. \tag{47}$$

Notice that this means that $\eta_\mathcal{H}$ depends on k. Hence there are three regions of interest, which are denoted by their conventional names and treated in the following way:

- **sub-horizon ($\mathcal{H} \ll k$)**: This region is irrelevant for the computation of the correlators at the end of inflation (see (46)), since solving (40) with the $\mathcal{H}^2\tilde{\Omega}$ term neglected with respect to the k^2 term we find
$$Q(\eta) = \frac{1}{\sqrt{2k}}\, R(\eta)\, e^{ik(\eta-\eta_i)} U \qquad \Rightarrow \qquad Q^*(\eta) Q^T(\eta) = \frac{1}{2k}\,\mathbb{1}. \tag{48}$$
(Here the normalization is fixed by the initial condition (45).) The end of the sub-horizon period η_- is therefore defined as the moment when this does not hold any more to first order, leading to the definition $\mathcal{H}^2(\eta_-) = \tilde{\epsilon}^{3/2} k^2$.‖
- **transition ($\mathcal{H} \sim k$)**: We consider (40) for Q, keeping all terms, but using that for a sufficiently small interval around $\eta_\mathcal{H}$ the slow-roll functions can be taken to be constant to first order, which makes it possible to obtain solutions for Q valid

---

§ There could be a problem with this approach, because our knowledge of physics beyond the Planck scale is extremely poor. In particular, the dispersion relation $\omega(\boldsymbol{k}) = k$ that we used implicitly might not be valid for large k: there might be a cut-off for large momenta. For a discussion of this trans-Planckian problem and possible cosmological consequences see [43, 44].

‖ The value 3/2 is chosen here because that is the same order to which the slow-roll background field equation is valid, see (14), but the arguments are independent of which specific power (larger than one) is chosen.

to first order using Hankel functions. Since the effect of the sub-horizon region is irrelevant, we take the following initial conditions:

$$Q(\eta_-) = \frac{1}{\sqrt{2k}}\,\mathbb{1}, \qquad Q'(\eta_-) = \frac{i\sqrt{k}}{\sqrt{2}}\,\mathbb{1}, \qquad R(\eta_-) = \mathbb{1}. \tag{49}$$

- **super-horizon ($\mathcal{H} \gg k$)**: In this region we use u to compute the vacuum correlator of the Newtonian potential Φ, which is related to u via a simple rescaling, see (27). As the k^2 dependence can be neglected, the exact solution for u of equation (34) is

$$u_{\boldsymbol{k}}(\eta) = u_{P\,\boldsymbol{k}} + C_{\boldsymbol{k}}\theta + D_{\boldsymbol{k}}\theta \int_{\eta_{\mathcal{H}}}^{\eta} \frac{\mathrm{d}\eta'}{\theta^2(\eta')}, \quad u_{P\,\boldsymbol{k}} = \theta \int_{\eta_{\mathcal{H}}}^{\eta} \frac{\mathrm{d}\eta'}{\theta^2} \int_{\eta_{\mathcal{H}}}^{\eta'} \mathrm{d}\eta''\, \mathcal{H}\theta\tilde{\eta}^{\perp} q_{2\,\boldsymbol{k}}, \tag{50}$$

  with $C_{\boldsymbol{k}}$ and $D_{\boldsymbol{k}}$ integration constants and $u_{P\,\boldsymbol{k}}$ a particular solution. To work out u_P in a more explicit form and to find solutions for Q slow-roll assumptions are necessary, which are treated in section 3.4.

As the sub-horizon region is irrelevant, what remains is the connection between the transition and the super-horizon region. In both these regions we have constructed analytic solutions of the same differential equation for Q. The only thing that must still be computed to determine the super-horizon solution uniquely, is the relative overall normalization between the solutions in these two regions. Instead of the more standard continuously differentiable matching at a specific time scale, we do this by identifying leading order asymptotic expansions.

This procedure works as follows. We can write both these solutions as power series in $k\eta$ and compare them in the transition region. There we find that the leading powers of the transition and super-horizon solutions are the same, separately for both the decaying and the non-decaying independent solution. The ratio of the coefficients in front of these leading powers gives us the relative normalization of the super-horizon solution with respect to the transition (and sub-horizon) solution. Although we need to compute the coefficients accurately to first order in slow roll, zeroth order turns out to be sufficient to distinguish the two independent solutions and identify the exponents of the leading terms in the expansions, see below (64). To conclude, we can determine the solution valid in the super-horizon region uniquely from the solution in the transition region around $\eta_{\mathcal{H}}$, even though the solution in the region in between is only known asymptotically. Some remarks on other matching schemes can be found at the end of section 3.5.

3.4. Slow roll for the perturbations

To determine the solution for Q in the super-horizon region, and to rewrite the particular solution u_P in terms of background quantities only, the concept of slow roll on the perturbations is useful. We now justify the use of slow roll on the perturbations and make this notion more precise. Physically it represents the fact that the combination of background and perturbation modes far outside the horizon cannot be distinguished from the background. We introduce the substitutions

$$\boldsymbol{\phi} \rightarrow \tilde{\boldsymbol{\phi}} = \boldsymbol{\phi} + \boldsymbol{\delta\phi}, \qquad b \rightarrow \tilde{b} = a(1+\Phi), \qquad a \rightarrow \tilde{a} = a(1-\Phi), \tag{51}$$

where we have chosen to work with conformal time after substitution to make a direct comparison with section 3.1 possible. Notice that in this way the perturbed metric (20) is obtained. Applying these substitutions to (6) and linearizing gives the perturbation equation (23) with k^2 put to zero, including the field curvature term. At the same time, by linearizing the combination

$$\mathcal{D}H_a + 3H_a^2 - \kappa^2 b^2 V = 0 \tag{52}$$

of the Friedmann equations (7), the equation of motion (21) for Φ is obtained. In other words, for the super-horizon modes the system of background equations (14) and (52) for $(\boldsymbol{\phi}, a, b)$ is also valid for the perturbed fields $(\tilde{\boldsymbol{\phi}}, \tilde{a}, \tilde{b})$. Hence the solutions for $(\boldsymbol{\phi}, a, b)$ and $(\tilde{\boldsymbol{\phi}}, \tilde{a}, \tilde{b})$ can only differ in their initial conditions, so that the perturbation quantities $(\boldsymbol{\delta\phi}, \Phi)$ are obtained by linearizing the background quantities with respect to the initial conditions:

$$\boldsymbol{\delta\phi} = (\boldsymbol{\nabla}_{\phi_0}\boldsymbol{\phi})\boldsymbol{\delta\phi}_0, \qquad\qquad \mathbf{P}^{\perp}\boldsymbol{q} = a\,\mathbf{P}^{\perp}\boldsymbol{\delta\phi}. \tag{53}$$

This technique was also used in [47, 48]. Here we have set the variations of the initial conditions a_0 and b_0 equal to zero, as a simple counting argument shows that this is sufficient to generate a complete set of solutions. Now if slow roll is valid for the background, it follows immediately that slow roll also governs the super-horizon perturbations. This fact has been used previously in the literature, see e.g. [23, 22].

Applying slow roll to the equation of motion (40) and rewriting it in terms of the quantity $Q_{SR} \equiv \frac{a_{\mathcal{H}}\sqrt{\tilde{\epsilon}_{\mathcal{H}}}}{a\sqrt{\tilde{\epsilon}}}\, Q Q_{\mathcal{H}}^{-1}$, we find

$$Q'_{SR} + \mathcal{H}\left[-\delta + (2\tilde{\epsilon} + \tilde{\eta}^{\|})\mathbb{1} + Z\right] Q_{SR} = 0, \qquad\qquad Q_{SR}(\eta_{\mathcal{H}}) = \mathbb{1}. \tag{54}$$

Here we have used that $\mathcal{D}\tilde{Q} = \tilde{Q}' - \mathcal{H}\tilde{Q}$ and $\mathcal{D}^2\tilde{Q} = \tilde{Q}'' - 3\mathcal{H}\tilde{Q}' - (\mathcal{H}' - 2\mathcal{H}^2)\tilde{Q}$, because $\tilde{Q}$ scales with one power of a. For reasons that will become clear in the next section we have defined δ as

$$\delta = -\frac{1}{3}\left(2\mathbb{1} + \frac{\Omega}{(1-\tilde{\epsilon})^2}\right) = \tilde{\epsilon}\,\mathbb{1} - \frac{\tilde{M}^2}{3H^2} + 2\tilde{\epsilon}\, e_1 e_1^T, \tag{55}$$

where the second expression is valid to first order. We make the additional assumption that also those components of $\tilde{M}^2/H^2$ that cannot be expressed in terms of the slow-roll functions defined in (10) are of first order, so that δ is a first-order quantity. The solution of (54) is found by integrating:

$$Q_{SR}(\eta) = \exp\left[\int_{\eta_{\mathcal{H}}}^{\eta} \mathrm{d}\eta'\, \mathcal{H}\left(\delta - (2\tilde{\epsilon} + \tilde{\eta}^{\|})\mathbb{1} - Z\right)\right]. \tag{56}$$

Although the initial conditions are applied at $\eta_{\mathcal{H}}$, this solution is only valid in the super-horizon region because k^2 terms have been neglected. Since slow roll has been used, this result is a priori not expected to be very accurate at the end of inflation. However, using the fact that the matrix between the brackets in (54) has its $(m1)$ components $(m \geq 1)$ all equal to zero to first order in slow roll (see (31) and (19)), one can easily find the solutions $(Q_{SR})_{11} = 1$ and $(Q_{SR})_{n1} = 0$ $(n > 1)$. Since these vary slowly (or rather not at all) even at the end of inflation, slow roll is still a good approximation there for these components.

3.5. Solutions of the perturbation equations to first order: calculation

In this section we perform the calculation that was discussed in section 3.3. As mentioned there, in a sufficiently small interval around $\eta_\mathcal{H}$ in the transition region the slow-roll functions can be taken constant. With this approximation we can obtain an expression for $\mathcal{H}(\eta)$ by integrating the relation for $\mathcal{H}'$ in (33) with respect to conformal time, while integrating $N' = \mathcal{H}$ gives the number of e-folds to first order around $\eta = \eta_\mathcal{H}$:

$$\mathcal{H}(\eta) = \frac{-1}{(1-\tilde{\epsilon}_\mathcal{H})\eta}, \qquad N(\eta) = N_\mathcal{H} - \frac{1}{1-\tilde{\epsilon}_\mathcal{H}} \ln\frac{\eta}{\eta_\mathcal{H}}. \tag{57}$$

Here we used the freedom in the definition of conformal time to set $\eta_\mathcal{H} = -1/[(1-\tilde{\epsilon}_\mathcal{H})k]$. From (32) we infer that to first order around $\eta = \eta_\mathcal{H}$

$$\theta(z) = \theta_\mathcal{H}\left(\frac{z}{z_\mathcal{H}}\right)^{1+2\tilde{\epsilon}_\mathcal{H}+\tilde{\eta}^\parallel_\mathcal{H}}, \qquad z \equiv k\eta, \qquad \theta_\mathcal{H} = \frac{\kappa}{\sqrt{2}}\frac{H_\mathcal{H}}{k\sqrt{\tilde{\epsilon}_\mathcal{H}}}. \tag{58}$$

In these expressions we have made the conventional choice of $\eta_\mathcal{H}$ as reference time to compute the constant slow-roll functions, etc., although in principle one could do the complete computation with another reference time scale. However, to be able to take $\tilde{\Omega}$ constant (see below), this time should not be much later than $\eta_\mathcal{H}$.

With the initial condition (49) the solution of (40) for the rotation matrix R during the transition region is

$$R(z) = e^{(N-N_-)Z_\mathcal{H}} = \left(\frac{z}{z_-}\right)^{-\frac{1}{1-\tilde{\epsilon}_\mathcal{H}}Z_\mathcal{H}}. \tag{59}$$

The only time dependent terms in the matrix Ω (30) are first order, so that we can take $\Omega = \Omega_\mathcal{H}$ in the transition region. The matrix $\tilde{\Omega}$ (40) on the other hand is given by

$$\tilde{\Omega} = R^{-1}(z)\Omega_\mathcal{H}R(z) = \Omega_\mathcal{H} - [\Omega_\mathcal{H}, Z_\mathcal{H}]\ln\frac{z}{z_-} = \Omega_\mathcal{H} + 3[\delta_\mathcal{H}, Z_\mathcal{H}]\left(\ln\frac{z}{z_\mathcal{H}} + \frac{3}{4}\ln\tilde{\epsilon}_\mathcal{H}\right), \tag{60}$$

where we used the definition of η_- from the previous section and $\delta_\mathcal{H} = \delta(\eta_\mathcal{H})$ is defined in (55). In this section we are still considering a single, arbitrary mode k (see equation (47) and the text above equation (35)). However, in the end we are interested in those modes that are visible in the CMBR, which crossed the Hubble scale in a small interval about 60 e-folds before the end of inflation. For those modes we estimate $\tilde{\epsilon}_\mathcal{H} \sim 0.01$ (motivated for example by a quadratic potential, see (83)), so that $\ln\tilde{\epsilon}_\mathcal{H} \sim \tilde{\epsilon}_\mathcal{H}^{-1/2}$. Since both δ and Z are of first order, the time dependence of $\tilde{\Omega}$ caused by the rotation is then only important at order 3/2 in the region around $z_\mathcal{H}$. (For a smaller value of $\tilde{\epsilon}_\mathcal{H}$ the effect is even of higher order.) Hence we take $\tilde{\Omega} = \Omega_\mathcal{H}$. From the correction term in equation (60) we can always check explicitly if that assumption is justified.

For matching in the region around $z_\mathcal{H}$ it will be useful to define $\bar{Q}(z) \equiv R(z_\mathcal{H})\tilde{Q}(z)$. Then $Q(z) = \bar{Q}(z)$ to first order in a sufficiently small region around $z_\mathcal{H}$. Using the same argument as in (60) the corresponding $\bar{\Omega} \equiv R(z_\mathcal{H})\tilde{\Omega}R^{-1}(z_\mathcal{H})$ is equal to $\Omega_\mathcal{H}$ to first order. Using this result and equation (57) for $\mathcal{H}$, equation (40) for $\tilde{Q}$ can be rewritten as an equation for $\bar{Q}$:

$$\bar{Q}_{,zz} + \bar{Q} - \frac{\nu_\mathcal{H}^2 - \frac{1}{4}}{z^2}\bar{Q} = 0, \qquad \nu_\mathcal{H}^2 = \frac{9}{4}\mathbb{1} + 3\delta_\mathcal{H}. \tag{61}$$

The solution of this matrix equation can be written in terms of a Hankel function:¶

$$\bar{Q}(z) = \sqrt{\frac{\pi}{4k}}\,\sqrt{z}H^{(1)}_{\nu_{\mathcal{H}}}(z), \qquad\qquad \nu_{\mathcal{H}} = \frac{3}{2}\mathbb{1} + \delta_{\mathcal{H}}. \tag{62}$$

Here the initial conditions (49) at the beginning of the transition region have been taken into account, as can be seen by using the fact that for $|z| \gg 1$ the Hankel function can be approximated by $H^{(1)}_{\nu}(z) = \sqrt{2/(\pi z)}\exp i(z - \pi\nu/2 - \pi/4)$ and neglecting unitary matrices. We also need the leading order term in the expansion in z of this result for $\bar{Q}$:

$$\bar{Q}_{lo} = \frac{1}{i\sqrt{2\pi k}}\,\Gamma(\nu_{\mathcal{H}})\left(\frac{z}{2}\right)^{\frac{1}{2}\mathbb{1}-\nu_{\mathcal{H}}} = -\frac{e^{-i\pi\delta_{\mathcal{H}}}}{i\sqrt{2k}}\,E_{\mathcal{H}}\left(\frac{z}{z_{\mathcal{H}}}\right)^{-\mathbb{1}-\delta_{\mathcal{H}}} \tag{63}$$

with

$$E_{\mathcal{H}} \equiv (1 - \tilde{\epsilon}_{\mathcal{H}})\mathbb{1} + (2 - \gamma - \ln 2)\delta_{\mathcal{H}}, \tag{64}$$

where $\gamma \approx 0.5772$ is the Euler constant. For later convenience we have defined the matrix $E_{\mathcal{H}}$, which to zeroth order in slow roll is equal to the identity. In (63) we have taken only the growing solution; the decaying one starts off with a term proportional to $z^{\frac{1}{2}\mathbb{1}+\nu_{\mathcal{H}}} = z^{+2\mathbb{1}+\delta_{\mathcal{H}}}$. We see that these two solutions can be distinguished already at zeroth order.

Next we turn to the super-horizon region. Here we have to relate u and Q by means of the first equation of (35). The solution for u is given in (50), while for q_1 we derive the following slow-roll equation of motion (see section 3.4):

$$q_1' - \frac{(1/\theta)'}{1/\theta}\,q_1 = 2\mathcal{H}\tilde{\eta}^{\perp}q_2 \quad\Rightarrow\quad q_1 = d\,\frac{1}{\theta} + 2\frac{1}{\theta}\int_{\eta_{\mathcal{H}}}^{\eta} \mathrm{d}\eta'\,\mathcal{H}\theta\tilde{\eta}^{\perp}q_2, \tag{65}$$

where we also gave the solution. By using slow roll we have selected the non-decaying solution for q_1/a. Using (35) we then find that the integration constant $D_{\boldsymbol{k}}$ in the solution (50) for u is given by $D_{\boldsymbol{k}} = \frac{1}{2}d$. The constant $C_{\boldsymbol{k}}$ is irrelevant because the function θ rapidly decays. The integration constant d can be determined using the procedure of identification of leading order terms (leading order in the expansion in z, not slow roll) described in section 3.3. Extrapolating the super-horizon solution for q_1 into the transition region sufficiently close to $\eta_{\mathcal{H}}$ that the integral can be neglected and that $\delta_{\mathcal{H}}\ln(z/z_{\mathcal{H}})$ is smaller than first order, and using (58) we find to first order $q_1 = (d/\theta_{\mathcal{H}})(z/z_{\mathcal{H}})^{-1}$. Under these conditions $e_1^T E_{\mathcal{H}}(z/z_{\mathcal{H}})^{-\mathbb{1}-\delta_{\mathcal{H}}} = e_1^T E_{\mathcal{H}}(z/z_{\mathcal{H}})^{-1}$ so that we can determine the constant d from equation (63). (Notice that, as mentioned in section 3.3, the exponents of $z/z_{\mathcal{H}}$ need only be identified to zeroth order, so that strictly speaking the condition that $\delta_{\mathcal{H}}\ln(z/z_{\mathcal{H}})$ is smaller than first order is not even necessary.) The final first-order result for $D_{\boldsymbol{k}}$ is:

$$\hat{D}_{\boldsymbol{k}} = -\frac{1}{2}\frac{e^{-i\pi\delta_{\mathcal{H}}}}{i\sqrt{2k}}\,\theta_{\mathcal{H}}e_1^T E_{\mathcal{H}}\hat{a}^{\dagger} + \text{c.c.}. \tag{66}$$

¶ This Bessel equation and its solution in terms of Hankel functions are well-known in the theory of inflationary density perturbations, see e.g. [2, 9] and references therein. However, in the multiple field case under consideration the order ν of the Hankel function is matrix valued. This should be considered in the usual way: defined by means of a series expansion.

For later use we note that this identification procedure can also be used for the complete matrix Q, not just for q_1. Completely analogously to (56) one has $\bar{Q} \propto a \exp\left(\int_{\eta_\mathcal{H}}^{\eta} \mathrm{d}\eta' \, \mathcal{H} \, R(\eta_\mathcal{H}) R^{-1} \left[\delta - \tilde{\epsilon} \mathbb{1}\right] R R^{-1}(\eta_\mathcal{H})\right)$, which behaves like $z^{-\mathbb{1}-\delta_\mathcal{H}}$ when extrapolated into the transition region around $\eta_\mathcal{H}$ using (57) and (60). Comparing with (63) we see that it is exactly the leading order term in the expansion of the Hankel function in the solution for Q in the transition region that goes over into the dominant solution for Q in the super-horizon region.

With this and (27) we can give the Newtonian potential Φ as a quantum operator at late times during inflation up to first order in slow roll:

$$\hat{\Phi}_{\boldsymbol{k}}(t) = -\frac{\kappa}{2ik^{3/2}} \frac{H_\mathcal{H}}{\sqrt{\tilde{\epsilon}_\mathcal{H}}} \left(A(t_\mathcal{H}, t) \, e_1^T + \tilde{U}_P^T(t)\right) E_\mathcal{H} e^{-i\pi\delta_\mathcal{H}} \hat{a}_{\boldsymbol{k}}^\dagger + \text{c.c.}. \tag{67}$$

Here we used the identity $\mathcal{H}_\mathcal{H} = k$ and the functions $A(t_\mathcal{H}, t)$ and $\tilde{U}_P^T(t)$ are defined as

$$A(t_\mathcal{H}, t) = 1 - \frac{H(t)}{a(t)} \int_{t_\mathcal{H}}^{t} \mathrm{d}t' a(t'), \qquad \tilde{U}_P^T = \frac{H}{a} \int_{\eta_\mathcal{H}}^{\eta} \mathrm{d}\eta' a^2 \tilde{\epsilon} \, U_P^T(\eta'),$$
$$U_P^T = 2\sqrt{\tilde{\epsilon}_\mathcal{H}} \int_{\eta_\mathcal{H}}^{\eta} \mathrm{d}\eta' \, \mathcal{H} \frac{\tilde{\eta}^\perp}{\sqrt{\tilde{\epsilon}}} \frac{a_\mathcal{H}}{a} \, e_2^T Q Q_\mathcal{H}^{-1}. \tag{68}$$

In $A(t_\mathcal{H}, t)$ we neglected one term which is exponentially suppressed with the number of e-folds. In this and all following equations $Q_\mathcal{H}$ is defined as the leading order asymptotic expression for Q evaluated at $\eta_\mathcal{H}$, i.e. $Q_\mathcal{H} = -E_\mathcal{H} \exp(-i\pi\delta_\mathcal{H})/(i\sqrt{2k})$. Remember that $\Phi_{\boldsymbol{k}}$ depends on k not only explicitly, but also implicitly through the dependence on $\eta_\mathcal{H}$.

Using slow roll on the perturbations and substituting the result for Q_{SR} from (56) into the definition for U_P^T we find

$$U_P^T = 2 \int_{\eta_\mathcal{H}}^{\eta} \mathrm{d}\eta' \, \mathcal{H} \tilde{\eta}^\perp \, e_2^T \exp\left[\int_{\eta_\mathcal{H}}^{\eta'} \mathrm{d}\eta'' \, \mathcal{H} \left(\delta - (2\tilde{\epsilon} + \tilde{\eta}^\parallel)\mathbb{1} - Z\right)\right] \tag{69}$$

to first order in slow roll. This expression is given in terms of background quantities only. U_P has no component in the e_1 direction, since to first order $(Q_{SR})_{21} = 0$. In section 4 we show how U_P^T can be computed explicitly for the case of a quadratic potential on a flat field manifold using the concept of slow roll on the perturbations.

We have been able to determine the integration constant $D_{\boldsymbol{k}}$ in the solution for u in the super-horizon region to first order in slow roll by using analytic properties of the solutions for Q in the transition region. We did not have to resort to a continuously differentiable matching at a specific time scale; the only time scale that appears in the result is the reference time $\eta_\mathcal{H}$ in the neighbourhood of which we have expanded the solutions. In the literature the concept of matching at a specific time is often used (see e.g. [6, 9]), for which usually the time of horizon crossing of either a generic or specific mode k is used. On the one hand matching for the scales of observational interest is then performed at times when $k\eta \approx 1$, while on the other hand approximations only valid for small $k\eta$ are used. The identification procedure described in section 3.3 and used in this section shows why the standard (single-field) results in the literature are nonetheless correct: by neglecting the k dependent corrections and taking $k\eta = 1$ (i.e.

$z/z_{\mathcal{H}} = 1$) one is exactly computing the overall normalization factor that we showed to be the only thing that needs to be determined. (Another possible way to solve the problem would be to match the transition and super-horizon solutions at a specific time η_+ later than $\eta_{\mathcal{H}}$, so that $k\eta_+ \ll 1$ is a valid assumption to first order. However, it turns out that the interval between $\eta_{\mathcal{H}}$ and this η_+ is too large to satisfy the requirement that the slow-roll functions can be taken constant, so that the Hankel solutions are not valid over the whole interval to first order.)

3.6. Adiabatic and isocurvature perturbations after inflation

The main subject of this paper is the treatment of perturbations during inflation, culminating in the result (67) for Φ for super-horizon modes during inflation. (The result for the other perturbations Q is given by (56) with $Q_{\mathcal{H}} = -E_{\mathcal{H}} \exp(-i\pi\delta_{\mathcal{H}})/(i\sqrt{2k})$.) However, in this section we discuss the relation between these results and the relevant quantities at the time of recombination when the CMBR was formed. Regarding the isocurvature perturbations after inflation we follow the treatment by Langlois [24], making several simplifying assumptions. Other work on this subject can be found in e.g. [49, 23, 50, 18, 51] and references therein.

Conventionally (see e.g. [18]) perturbations are divided into adiabatic (or curvature) and entropy (or isocurvature) perturbations. Adiabatic perturbations are perturbations in the total energy density, and are the only ones present in the case of single-field inflation. If there are N scalar fields, there are in general $N - 1$ isocurvature perturbations in addition to the single adiabatic perturbation. Isocurvature perturbations are perturbations in the ratios of energy densities of the different components, leaving the total energy density unperturbed. With our basis (8) the adiabatic perturbation corresponds with the e_1 component, while the isocurvature perturbations correspond in principle with all the other components. These different types of perturbations can be sources for each other, see e.g. equation (34). By including the particular solution in the result for Φ (67) we have taken into account the effect of the isocurvature perturbations on the adiabatic perturbation during inflation.

Although the treatment of purely adiabatic perturbations after inflation is rather straightforward, this is not the case for isocurvature perturbations. The evolution of the isocurvature perturbations depends on some additional physical input, most importantly to what kind of particles the multiple scalar fields decay. If the interactions between these different kinds of particles are too strong, the isocurvature perturbations might not survive till the time of recombination at all [24]. We assume that one of the fields decays to all the standard model particles, while the other fields decay to different kinds of cold dark matter, so that there is no interaction between the decay products of the different fields (except gravitational) and they can be considered as ideal fluid components without mutual interactions. We also neglect the effects of (p)reheating and assume that at the end of inflation there is an immediate transition to a radiation dominated universe. For purely adiabatic perturbations the presence of preheating is irrelevant on super-horizon scales, as was proved in [52]. In the presence of isocurvature perturbations preheating may have an important effect on the perturbations (see e.g. [53, 54] and references therein), but that is beyond the scope of this paper.

We first give the three vacuum correlators of the gravitational potential valid up to and including first order in slow roll at the time of recombination, and discuss their derivations afterwards. These three are the adiabatic contribution (but including isocurvature effects during inflation), the isocurvature contribution, and the mixing between them:

$$\begin{aligned}\langle\Phi^2_{\boldsymbol{k}\,ad}\rangle_{t_{rec}} \quad &= \frac{9}{25}\,\frac{\kappa^2}{4k^3}\frac{H^2_{\mathcal{H}}}{\tilde{\epsilon}_{\mathcal{H}}}\Big[(1-2\tilde{\epsilon}_{\mathcal{H}})(1+U^T_{P\,e}U_{P\,e}) \\ &\qquad + 2B\left((2\tilde{\epsilon}_{\mathcal{H}}+\tilde{\eta}^{\parallel}_{\mathcal{H}}) + 2\tilde{\eta}^{\perp}_{\mathcal{H}}e_2^T U_{P\,e} + U^T_{P\,e}\delta_{\mathcal{H}}U_{P\,e}\right)\Big],\end{aligned} \tag{70}$$

$$\langle\Phi^2_{\boldsymbol{k}\,iso}\rangle_{t_{rec}} \quad = \frac{1}{25}\,\frac{\kappa^2}{4k^3}\frac{H^2_{\mathcal{H}}}{\tilde{\epsilon}_{\mathcal{H}}}\Big[(1-2\tilde{\epsilon}_{\mathcal{H}})V_e^T V_e + 2BV_e^T\delta_{\mathcal{H}}V_e\Big], \tag{71}$$

$$\langle\Phi_{\boldsymbol{k}\,iso}\Phi_{\boldsymbol{k}\,ad}\rangle = \frac{3}{25}\,\frac{\kappa^2}{4k^3}\frac{H^2_{\mathcal{H}}}{\tilde{\epsilon}_{\mathcal{H}}}\Big[(1-2\tilde{\epsilon}_{\mathcal{H}})V_e^T U_{P\,e} + 2B\left(\tilde{\eta}^{\perp}_{\mathcal{H}}e_2^T V_e + V_e^T\delta_{\mathcal{H}}U_{P\,e}\right)\Big], \tag{72}$$

with $U^T_{P\,e} \equiv U^T_P(\eta_e)$ given in (68) and V_e defined by

$$V_e^T = \frac{1}{2}\sqrt{\tilde{\epsilon}_{\mathcal{H}}}\,\frac{\sqrt{\tilde{\epsilon}_e}\,\tilde{\eta}^{\perp}_e}{\tilde{\epsilon}_e+\tilde{\eta}^{\parallel}_e}\frac{a_{\mathcal{H}}}{a_e}\,e_2^T Q_e Q^{-1}_{\mathcal{H}}. \tag{73}$$

Here $B \equiv 2-\gamma-\ln 2 \approx 0.7297$ and we used the definition of $E_{\mathcal{H}}$ in (64) and the fact that $U_{P\,e}$ and V_e have no components in the e_1 direction since $(a_{\mathcal{H}}/a)(QQ^{-1}_{\mathcal{H}})_{21} = 0$, see below (56). Moreover, we assume V_e to be of order 1 at most, otherwise other terms have to be included in the vacuum correlators to give the complete results to first order in slow roll.

The explicit multiple field terms are the contributions of the terms $U_{P\,e}$ and V_e, which are absent in the single-field case. Since they are both to a large extent determined by $\tilde{\eta}^{\perp}$, we see that the behaviour of $\tilde{\eta}^{\perp}$ during the last 60 e-folds of inflation is crucial to determine whether multiple-field effects are important. For example, one can immediately draw the conclusion that in assisted inflation [55], where one quickly goes to an attractor solution with all $\dot{\phi}_i$ equal to each other apart from constant factors, so that $\tilde{\eta}^{\perp} = 0$, there will be no explicit multiple-field contributions to the gravitational potential. One can draw the conclusion that with our basis the total isocurvature perturbation that appears in the expressions for Φ only depends on the e_2 component of q, independently of the total number of fields and the actual number of independent entropy perturbations. The fact that the entropy perturbations act as sources for the adiabatic perturbation (the U_P term) naturally leads to correlations between adiabatic and isocurvature perturbations, as described by the mixed correlator. In fact these correlations are only absent if the source term disappears, i.e. again if $\tilde{\eta}^{\perp}$ vanishes. (The authors of [18] studied the two-field case and found the derivative of the angle that parameterizes the influence of the second field on the background trajectory to be the relevant parameter. In the two-field limit this parameter corresponds with $\tilde{\eta}^{\perp}$, but our result is valid for an arbitrary number of fields.) In [24] these correlations were studied in the context of a double inflation model.

Using the concept of slow roll on the perturbations, introduced in section 3.4, the quantity $U_{P\,e}$ can be rewritten in terms of background quantities only, as was done in (69). Because we have used slow roll in the derivation, this expression is in principle not valid at the very end of inflation. If (69) does indeed give a bad approximation for $U_{P\,e}$, for example if $\tilde{\eta}^{\perp}$ grows very large, a more careful treatment of the transition at the end of inflation is necessary. However, in other cases the contribution to the integral near the end of inflation can be negligible, for example if $\tilde{\eta}^{\perp}$ goes sufficiently rapidly to zero at the end of inflation. In those cases (69) gives a very good approximation for $U_{P\,e}$ and the details of the transition at the end of inflation are unimportant for the gravitational correlator (70). An important example of this latter case is discussed in section 4. Unfortunately V_e depends very much on the details of the transition at the end of inflation, so that for an actual calculation a model of this transition has to be assumed, a treatment of which is beyond the scope of this paper.

We conclude this section by discussing the derivation of the correlators. Assuming the absence of anisotropic stress, the equation for super-horizon modes of Φ derived from the (00) and (ij) components of the Einstein equations can be written as (see e.g. [6, 26]):

$$\Phi'' + 3\mathcal{H}(1+c_s^2)\Phi' + \kappa^2 a^2(\rho c_s^2 - p)\Phi = \frac{1}{2}\kappa^2 a^2 e \tag{74}$$

with

$$\rho \equiv -T^0_{\ 0} = \frac{3}{\kappa^2 a^2}\mathcal{H}^2, \quad p \equiv \frac{1}{3}T^i_{\ i} = -\frac{1}{\kappa^2 a^2}(2\mathcal{H}' + \mathcal{H}^2), \quad c_s^2 \equiv \frac{p'}{\rho'}, \quad e \equiv \delta p - c_s^2 \delta\rho. \tag{75}$$

Here ρ and p are the energy density and isotropic pressure, c_s^2 is the sound velocity and e is the total (gauge invariant) entropy perturbation. The expressions after the second equality sign in the definitions of ρ and p follow from the background Einstein equations. The perturbations are analogously defined as $\delta\rho \equiv -\delta T^0_{\ 0}$ and $\delta p \equiv \frac{1}{3}\delta T^i_{\ i}$.

The adiabatic perturbation at the time of recombination is usually defined as the homogeneous solution of (74) at that time, where as initial conditions one has matched to the total solution for Φ at the beginning of the radiation dominated era. Hence the adiabatic perturbation does include the effects of isocurvature perturbations during inflation (and in general preheating). We can rewrite the homogeneous part of equation (74) as $u'' - (\theta''/\theta)u = 0$ with u and θ given by the same definitions (27) and (32), which are also defined after inflation in terms of a and its derivatives denoted by $\mathcal{H}$ and $\tilde{\epsilon}$. Hence we can use the same solution (50), with different constants $\tilde{C}_{\boldsymbol{k}}$ and $\tilde{D}_{\boldsymbol{k}}$ and without the particular solution. Matching our complete solution for Φ at the end of inflation t_e to the homogeneous solution for Φ after inflation, we find for the latter:

$$\begin{aligned} \Phi_{\boldsymbol{k}\,ad}(t) &= 2\left(D_{\boldsymbol{k}} + \theta_e\, u'_{P\,\boldsymbol{k}}(\eta_e) - \theta'_e\, u_{P\,\boldsymbol{k}}(\eta_e)\right) A(t_e,t) \\ &= -\frac{\kappa}{2ik^{3/2}}\frac{H_{\mathcal{H}}}{\sqrt{\tilde{\epsilon}_{\mathcal{H}}}} A(t_e,t)\left(e_1^T + U_{P\,e}^T\right) E_{\mathcal{H}} e^{-i\pi\delta_{\mathcal{H}}}\hat{a}^\dagger_{\boldsymbol{k}} + \text{c.c.} \end{aligned} \tag{76}$$

This expression is only valid some time after t_e, since we have neglected the $\tilde{C}_{\boldsymbol{k}}$ term, which is suppressed by H/a. The function A is defined in (68). At the time of recombination we can use that $a(t) \propto t^{2/3}$ in a matter dominated universe to find $A(t_e, t_{rec}) = 3/5$.[+]

The isocurvature contribution to the gravitational potential at the time of recombination is usually defined as the particular solution of (74), with the initial conditions that it is zero and has zero derivative at the beginning of the radiation dominated era. Following [24] we assume that $N-1$ fields have decayed to non-interacting cold dark matter (ideal fluid) components and the remaining field to the standard model particles, which we represent by the photons and denote by the subscript r. Then we can write (using [26]):

$$\tilde{S} \equiv -\frac{1}{4}\frac{e}{\rho c_s^2 - p} = \frac{\sum_{i=1}^{N-1} \rho_i S_i}{\sum_{i=1}^{N-1} \rho_i}, \tag{77}$$

where S_i is defined by $S_i \equiv \delta\rho_i/(\rho_i + p_i) - \delta\rho_r/(\rho_r + p_r)$ and we used $p_i = 0$ for the CDM components and $p_r = \rho_r/3$ for the photons. In [49, 26] it is proved that for the case of ideal fluid components without mutual interactions in a flat universe, the super-horizon modes of S_i are constant. Since the ρ_i all have the same time dependence $\propto a^{-3}$, this means that $\tilde{S}$ is also constant. Hence we find a very simple particular solution for Φ: $\Phi_p = -2\tilde{S}$. To take care of the initial conditions we have to add a part of the homogeneous solution and find

$$\Phi_{iso}(t) = -2\tilde{S} + 2\tilde{S}\left(1 + \frac{1}{\tilde{\epsilon}_e}\right) A(t_e, t). \tag{78}$$

Since by assumption $\tilde{\epsilon}_e = \tilde{\epsilon}_{rad} = 2$ and $A(t_e, t_{rec}) = 3/5$ we find at the time of recombination $\Phi_{iso}(t_{rec}) = -\tilde{S}/5$. To link $\tilde{S}$ to inflationary quantities one computes $\tilde{S}$ according to its definition (77) during inflation and makes the assumption that $\tilde{S}$(after inflation) $= \tilde{S}$(end of inflation). The result is:

$$\tilde{S} = \frac{\kappa}{2\sqrt{2}}\left[\frac{\sqrt{\tilde{\epsilon}}}{\tilde{\epsilon} + \tilde{\eta}^{\parallel}}\,\tilde{\eta}^{\perp}\,\frac{q_2}{a}\right]_e. \tag{79}$$

This is not a slow-roll approximated expression; the slow-roll functions are merely short-hand notation defined by (10).

4. Illustration: scalar fields with a quadratic potential

4.1. Analytical expressions for background and perturbations

In this section we consider slow-roll inflation with scalar fields living on the flat manifold $\mathbb{R}^N$ with a quadratic potential V. The slow-roll equation of motion and Friedmann

---

[+] Sometimes the curvature perturbation on super-horizon scales $-\zeta = \Phi - (H/\dot{H})(\dot{\Phi} + H\Phi)$ is used in the literature instead of the gravitational potential (see e.g. [18] and references therein), since it is constant for purely adiabatic perturbations [52]. However, as one can see from (76), this definition simply removes the time dependent factor A. Because of the expansion of the universe A has a well-defined value at recombination, independent of evolutionary details, so that one can use Φ as well.

equation for the background quantities to first order are given by

$$\dot{\boldsymbol{\phi}} = -\frac{2}{\sqrt{3}\,\kappa}\,\boldsymbol{\partial}^T\sqrt{V(\boldsymbol{\phi})}, \qquad H = \frac{\kappa}{\sqrt{3}}\,\sqrt{V(\boldsymbol{\phi})}\left(1+\frac{\tilde{\epsilon}}{6}\right), \qquad V = \frac{1}{2}\kappa^{-2}\,\boldsymbol{\phi}^T\mathbf{m}^2\boldsymbol{\phi}. \tag{80}$$

Here $\mathbf{m}^2$ is a general symmetric mass matrix given in units of the Planck mass κ^{-1}. The initial starting point of the field $\boldsymbol{\phi}$ is denoted by $\boldsymbol{\phi}_0 = \boldsymbol{\phi}(0)$. The solution of the equation of motion (80) can be written in terms of one dimensionless positive scalar function $\psi(t)$:

$$\boldsymbol{\phi}(t) = e^{-\frac{1}{2}\mathbf{m}^2\psi(t)}\boldsymbol{\phi}_0 \qquad \Rightarrow \qquad \dot{\psi} = \sqrt{\frac{2}{3}}\frac{2}{\kappa^2\phi_0}\left(\hat{\boldsymbol{\phi}}_0^T\,\mathbf{m}^2 e^{-\mathbf{m}^2\psi}\hat{\boldsymbol{\phi}}_0\right)^{-\frac{1}{2}}, \tag{81}$$

with the initial condition $\psi(0) = 0$ and where $\hat{\boldsymbol{\phi}}_0 \equiv \boldsymbol{\phi}_0/\phi_0$, with $\phi_0 \equiv |\boldsymbol{\phi}_0|$, denotes the unit vector in the direction of the initial position in field space. In other words, we have determined the trajectory that the field $\boldsymbol{\phi}$ follows through field space starting from point $\boldsymbol{\phi}_0$. The number of e-folds $N = \int H\mathrm{d}t$ and the slow-roll function $\tilde{\epsilon}$ can be given as a function of ψ by using (81):

$$\begin{aligned} N(\psi) &= N_\infty\left(1-\hat{\boldsymbol{\phi}}_0^T e^{-\mathbf{m}^2\psi}\hat{\boldsymbol{\phi}}_0\right) - \frac{1}{12}\ln\frac{\hat{\boldsymbol{\phi}}_0^T\,\mathbf{m}^2 e^{-\mathbf{m}^2\psi}\hat{\boldsymbol{\phi}}_0}{\hat{\boldsymbol{\phi}}_0^T\,\mathbf{m}^2\hat{\boldsymbol{\phi}}_0}, \\ \tilde{\epsilon}(\psi) &= \frac{1}{2N_\infty}\frac{\hat{\boldsymbol{\phi}}_0^T\,\mathbf{m}^4 e^{-\mathbf{m}^2\psi}\hat{\boldsymbol{\phi}}_0}{(\hat{\boldsymbol{\phi}}_0^T\,\mathbf{m}^2 e^{-\mathbf{m}^2\psi}\hat{\boldsymbol{\phi}}_0)^2} \end{aligned} \tag{82}$$

with $N_\infty = \frac{1}{4}\kappa^2\phi_0^2$. For the other slow-roll functions similar expressions can be obtained. The slow-roll limit for the total number of e-folds during inflation N_∞ is approached by taking the limit $\psi \to \infty$ in the zeroth order expression for N, i.e. the above expression without the logarithm.

It is useful to have a leading order estimate of $\tilde{\epsilon}_{\mathcal{H}}$. To this end we take the zeroth order expression for $N(\psi)$ and assume that ψ is already so large at time $t_{\mathcal{H}}$ that we can neglect all masses except the smallest one in the exponential $\exp(-\frac{1}{2}\mathbf{m}^2\psi)$. Then we can solve for $\psi_{\mathcal{H}}$ and insert this into the expression for $\tilde{\epsilon}(\psi)$ to find

$$e^{-m_1^2\psi_{\mathcal{H}}} = \frac{1}{||\mathbf{E}_1||^2}\frac{N_\infty - N_{\mathcal{H}}}{N_\infty} \qquad \Rightarrow \qquad \tilde{\epsilon}_{\mathcal{H}} = \frac{1}{2(N_\infty - N_{\mathcal{H}})}. \tag{83}$$

Here m_1 is the smallest mass eigenvalue, $\mathbf{E}_1$ is the projection operator that projects on the eigenspace of m_1 and $||\mathbf{E}_1||^2 = \hat{\boldsymbol{\phi}}_0{}^T\mathbf{E}_1\hat{\boldsymbol{\phi}}_0 \leq 1$. Since $N_\infty - N_{\mathcal{H}} \approx 60$ we see that $\tilde{\epsilon}_{\mathcal{H}} \sim 0.01$.

We continue by computing the particular solution $U_{P\,e} = U_P(\eta_e)$ defined in (68). It turns out that in this case we can work out the integral analytically in slow roll, making use of the fact that we have obtained the slow-roll trajectories in (81). The velocity and acceleration are given by

$$\dot{\boldsymbol{\phi}} = -\frac{1}{2}\dot{\psi}\,\mathbf{m}^2\boldsymbol{\phi}, \qquad \mathcal{D}_t\dot{\boldsymbol{\phi}} = -\frac{1}{2}\ddot{\psi}\,\mathbf{m}^2\boldsymbol{\phi} + \frac{1}{4}\dot{\psi}^2\,\mathbf{m}^4\boldsymbol{\phi}, \tag{84}$$

while according to (53) we obtain $\boldsymbol{\delta\phi}$ by varying $\boldsymbol{\phi}$ with respect to the initial conditions:

$$\boldsymbol{\delta\phi} = -\frac{1}{2}\delta\psi\,\mathbf{m}^2\boldsymbol{\phi} + e^{-\frac{1}{2}\mathbf{m}^2\psi}\boldsymbol{\delta\phi}_0, \tag{85}$$

where $\delta\psi$ is the function ψ varied with respect to $\boldsymbol{\phi}_0$. The projector parallel to the velocity is given by $\mathbf{P}^{\parallel} = \boldsymbol{\mathsf{m}}^2\boldsymbol{\phi}\,\boldsymbol{\phi}^T\boldsymbol{\mathsf{m}}^2/(\boldsymbol{\phi}^T\boldsymbol{\mathsf{m}}^4\boldsymbol{\phi})$, and therefore we find that

$$\mathcal{D}_t\dot{\boldsymbol{\phi}}^T\,\mathbf{P}^{\perp}\boldsymbol{\delta\phi} = \frac{1}{4}\dot{\psi}^2\,\boldsymbol{\phi}^T\left[\boldsymbol{\mathsf{m}}^4 - \frac{\boldsymbol{\phi}^T\boldsymbol{\mathsf{m}}^6\boldsymbol{\phi}}{\boldsymbol{\phi}^T\boldsymbol{\mathsf{m}}^4\boldsymbol{\phi}}\boldsymbol{\mathsf{m}}^2\right]e^{-\frac{1}{2}\boldsymbol{\mathsf{m}}^2\psi}\boldsymbol{\delta\phi}_0. \tag{86}$$

Here we have used that the first terms of $\mathcal{D}_t\dot{\boldsymbol{\phi}}$ and $\boldsymbol{\delta\phi}$ are proportional to $\dot{\boldsymbol{\phi}}$ and hence are projected away, so that $\delta\psi$ drops out. We rewrite $U^T_{P\,e}$ such that we can apply this result:

$$U^T_{P\,e}q_{\mathcal{H}} = 2\sqrt{\tilde{\epsilon}_{\mathcal{H}}}\int_{t_{\mathcal{H}}}^{t_e}\mathrm{d}t\,\frac{H}{\sqrt{\tilde{\epsilon}}}\,\tilde{\boldsymbol{\eta}}^T\mathbf{P}^{\perp}\,a_{\mathcal{H}}\boldsymbol{\delta\phi}. \tag{87}$$

Substituting the definition (10) for $\tilde{\boldsymbol{\eta}}$ and using (82) for $\tilde{\epsilon}$ and (84) to determine $|\dot{\boldsymbol{\phi}}|$, the integral takes the form

$$U^T_{P\,e}q_{\mathcal{H}} = \frac{\kappa\sqrt{\tilde{\epsilon}_{\mathcal{H}}}}{\sqrt{2}}\int_{\psi_{\mathcal{H}}}^{\psi_e}\mathrm{d}\psi\,\frac{\boldsymbol{\phi}^T\boldsymbol{\mathsf{m}}^2\boldsymbol{\phi}}{\boldsymbol{\phi}^T\boldsymbol{\mathsf{m}}^4\boldsymbol{\phi}}\,\boldsymbol{\phi}^T\boldsymbol{\mathsf{m}}^4\mathbf{P}^{\perp}e^{-\frac{1}{2}\boldsymbol{\mathsf{m}}^2(\psi-\psi_{\mathcal{H}})}\,a_{\mathcal{H}}\boldsymbol{\delta\phi}_{\mathcal{H}}. \tag{88}$$

By writing out the projector $\mathbf{P}^{\perp}$ we can employ

$$\frac{1}{\boldsymbol{\phi}^T\boldsymbol{\mathsf{m}}^4\boldsymbol{\phi}}\left[\boldsymbol{\phi}^T\boldsymbol{\mathsf{m}}^4 - \frac{\boldsymbol{\phi}^T\boldsymbol{\mathsf{m}}^6\boldsymbol{\phi}}{\boldsymbol{\phi}^T\boldsymbol{\mathsf{m}}^4\boldsymbol{\phi}}\boldsymbol{\phi}^T\boldsymbol{\mathsf{m}}^2\right]e^{-\frac{1}{2}\boldsymbol{\mathsf{m}}^2\psi} = -\frac{\partial}{\partial\psi}\left[\frac{\boldsymbol{\phi}^T\boldsymbol{\mathsf{m}}^2e^{-\frac{1}{2}\boldsymbol{\mathsf{m}}^2\psi}}{\boldsymbol{\phi}^T\boldsymbol{\mathsf{m}}^4\boldsymbol{\phi}}\right] \tag{89}$$

to perform a partial integration to express $U^T_{P\,e}$ as

$$U^T_{P\,e}q_{\mathcal{H}} = \frac{\kappa\sqrt{\tilde{\epsilon}_{\mathcal{H}}}}{\sqrt{2}}\left[\boldsymbol{\phi}^T\mathbf{P}^{\perp}e^{-\frac{1}{2}\boldsymbol{\mathsf{m}}^2(\psi-\psi_{\mathcal{H}})}\right]_{\psi_{\mathcal{H}}}^{\psi_e}\,a_{\mathcal{H}}\boldsymbol{\delta\phi}_{\mathcal{H}}. \tag{90}$$

To determine $a_{\mathcal{H}}\delta\phi_{\mathcal{H}}$ we use the definition of q in (27): $q_{\mathcal{H}} = a_{\mathcal{H}}(\delta\phi_{\mathcal{H}} + (\sqrt{2\tilde{\epsilon}_{\mathcal{H}}}/\kappa)\,\Phi_{\mathcal{H}}e_1)$, where we also inserted the definition of $\tilde{\epsilon}$. Using (27) and (36) to relate $\Phi_{\mathcal{H}}$ to $q_{\mathcal{H}}$ we obtain

$$a_{\mathcal{H}}\delta\phi_{\mathcal{H}} = X^Tq_{\mathcal{H}}, \qquad \text{with} \qquad X = 1 - \tilde{\epsilon}_{\mathcal{H}}\left(2\tilde{\eta}^{\perp}_{\mathcal{H}}e_2 + \left(2\tilde{\epsilon}_{\mathcal{H}} + \tilde{\eta}^{\parallel}_{\mathcal{H}} - \delta_{\mathcal{H}}\right)e_1\right)e_1^T, \tag{91}$$

where we made use of the relation $Q'_{\mathcal{H}} = \mathcal{H}_{\mathcal{H}}(1 - \tilde{\epsilon}_{\mathcal{H}} + \delta_{\mathcal{H}})Q_{\mathcal{H}}$ that follows from (63). Hence $X = 1$ to first order in slow roll. With this we find our final result for $U^T_{P\,e}$:

$$U^T_{P\,e} = \frac{\kappa\sqrt{\tilde{\epsilon}_{\mathcal{H}}}}{\sqrt{2}}\left(-\phi^{\perp}_{\mathcal{H}} + e^{-\frac{1}{2}\mathsf{m}^2(\psi_e-\psi_{\mathcal{H}})}\phi^{\perp}_e\right)^T. \tag{92}$$

Here all terms are written in terms of the basis $\{\boldsymbol{e}_n\}$: $\phi^{\perp}$ denotes the vector with components $\boldsymbol{e}_n^{\dagger}\mathbf{P}^{\perp}\boldsymbol{\phi}$ and $\exp(-\mathsf{m}^2\psi)$ the matrix with components $\boldsymbol{e}_m^{\dagger}\exp(-\boldsymbol{\mathsf{m}}^2\psi)\boldsymbol{e}_n$. The second term within the parentheses in the expression for $U^T_{P\,e}$ is in general very small. In the first place all but the least massive field will have reached zero near the end of inflation, so that $\phi^{\perp}_e$ is small. In the second place this term is suppressed by the large negative exponential, since ψ_e is very large near the end of inflation, even though we may not be able to take the limit of $\psi_e \to \infty$ since slow roll is then not valid anymore.



Figure 1. a) Background fields and b) slow-roll functions as a function of the number of e-folds in the model with two fields on a flat manifold with a quadratic potential with masses $m_1 = 1 \cdot 10^{-5}$, $m_2 = 2.5 \cdot 10^{-5}$ and initial conditions $\phi_1 = 20$, $\phi_2 = 25$.

4.2. Numerical example

We now treat a numerical example, not only to illustrate the theory, but also to check our analytical results. We take the situation of two fields, with masses $m_1 = 1 \cdot 10^{-5}$ and $m_2 = 2.5 \cdot 10^{-5}$ in units of the Planck mass. As initial conditions we choose $\phi_1 = 20$ and $\phi_2 = 25$, also in Planckian units. Then $N_\infty = 256.25$, while an exact numerical calculation gives a total amount of inflation of 257.8 e-folds before the oscillations start. We have chosen the overall normalization of the masses such that we get the correct order of magnitude for the amplitude of the density perturbations. Apart from giving sufficient inflation, the specific choice of initial conditions has no special meaning. We compute all background quantities exactly, as we want to check the accuracy of our analytical results for the perturbations. In figure 1 we have plotted the fields and slow-roll functions as a function of the number of e-folds. We see that the more massive field goes to zero more quickly than the less massive field, as expected from (81). Moreover, around the time that the second field reaches zero, all slow-roll functions show a bump. For the chosen masses and initial conditions the bumps are located during the last 60 e-folds. As mentioned in the paragraph below (73), for multiple-field effects to be important we need $\tilde{\eta}^{\perp}$ to be substantial during the last 60 e-folds. Hence this is a good model to look for multiple-field effects. Moreover, as we see from the figure, $\tilde{\eta}^{\perp}$ goes to zero at the end of inflation, so that we expect corrections to $U_{P\,e}$, caused by the break-down of slow roll at the end of inflation, to be small. Indeed, figure 2 shows that the contribution to $U_{P\,e}$ during the last few e-folds of inflation is negligible.

The results for the amplitude of the adiabatic vacuum correlator of the gravitational potential are summarized in table 1. We split the contributions to the correlator into a homogeneous part (all terms without $U_{P\,e}$) and a particular part (the rest, so including mixing terms). Everything is evaluated for the mode k that crossed the horizon 60 e-folds before the end of inflation. The last column gives the relative error between our first order analytical results (70) and (92) on the one hand, and the exact numerical



Figure 2. a) The particular contribution U_P to the gravitational correlator during the super-horizon region as a function of the number of e-folds. To show the relation with the behaviour of $\tilde{\eta}^\perp$, this slow-roll function has been plotted again in figure b), on the same horizontal scale as figure a).

Table 1. The amplitude of the adiabatic vacuum correlator of the gravitational potential $|\delta_{\boldsymbol{k}}|^2 = \frac{1}{2\pi^2} k^3 \langle \Phi^2_{\boldsymbol{k}\,ad}(t_{rec}) \rangle$ for the mode k that crossed the horizon 60 e-folds before the end of inflation, is separated into a purely homogeneous and a (mixed) particular part. The first two columns give their values and their relative contributions to the total correlator according to our analytical slow-roll result (70) combined with (92). The final column shows the relative error between these expressions and the exact numerical results.

	Amplitude $\lvert\delta_{\boldsymbol{k}}\rvert^2$	Contribution to total	Relative error
Homogeneous	$1.55 \cdot 10^{-9}$	0.505	0.0001
Particular	$1.52 \cdot 10^{-9}$	0.495	0.0006
Total	$3.08 \cdot 10^{-9}$	1	0.0003

result on the other. These results agree with our claim that we computed the correlator to first order in slow roll: the relative errors are (much) smaller than $\mathcal{O}(\tilde{\epsilon}_{\mathcal{H}})$. We also see that our slow-roll approximation for U_P is indeed still very good at the end of inflation. The column before that shows the relative contributions of the various parts to the total adiabatic correlator. We see that the particular solution terms are responsible for almost half the total result in this model. Hence neglecting these terms to leading order, which might naively be done because they couple with a $\tilde{\eta}^\perp$ in (34), can be dangerous.

For (significantly) larger or smaller mass ratios, $\tilde{\eta}^\perp$ is smaller during the last 60 e-folds and the contribution of the explicit multiple-field terms to the correlator is less important. This could also be expected a priori, since a much larger mass ratio means that the heavy field has already reached zero before the last 60 e-folds, and the situation is effectively single-field. On the other hand, a much smaller mass ratio means that we approach the limit of equal masses, which corresponds with a central potential that is also effectively single-field.

5. Conclusions and discussion

We have given a general treatment for scalar perturbations on a flat Robertson-Walker spacetime in the presence of an arbitrary number of scalar fields that take values on a curved field manifold during slow-roll inflation. These are the kind of systems that one typically obtains from (string-inspired) high-energy models. The scalar perturbations are calculated to first order in slow roll. In particular we compute the vacuum correlator of the gravitational potential in terms of background quantities only, which is related to the temperature fluctuations that are observed in the CMBR.

A discussion of the background scalar fields served as the foundation for this analysis. The first of three central ingredients for this discussion is the manifestly covariant treatment with respect to reparameterizations of the field manifold and of the time variable. Secondly, the field dynamics (the field velocity, acceleration, etc.) naturally induce an orthonormal basis $(\boldsymbol{e}_1, \boldsymbol{e}_2, \ldots)$ on the field manifold. This makes a separation between effectively single-field and truly multiple-field contributions possible. Finally, we modified the definitions of the well-known slow-roll parameters to define slow-roll functions in terms of derivatives of the Hubble parameter and the background field velocity for the case of multiple scalar field inflation. These slow-roll functions are vectors, which can be decomposed in the basis induced by the field dynamics. For example, the slow-roll function $\tilde{\eta}^{\perp}$ measures the size of the acceleration perpendicular to the field velocity. Because we did not make the assumption that slow roll is valid in the definition of the slow-roll functions, it is often possible to identify these slow-roll functions in exact equations of motion and make decisions about neglecting some of the terms. However, more important for precision calculations are estimates of the accuracy of the solutions of these approximated slow-roll equations; it turns out that if the size of the region of integration is too large this accuracy may be compromised.

Our calculation of the scalar perturbations accurate to first order in slow roll is based on the following cornerstones. We generalized the combined system of gravitational and matter perturbations of Mukhanov et al. [6] by defining the Mukhanov-Sasaki variables as a vector on the scalar field manifold. The decomposition of these variables in the basis induced by the background field dynamics is field space reparameterization invariant, and the corresponding Lagrangean takes the standard canonical form, making quantization straightforward. The gravitational potential only couples to the scalar field perturbation in the direction $\boldsymbol{e}_2$ with a slow-roll factor $\tilde{\eta}^{\perp}$.

To obtain analytic solutions for the scalar perturbations to first order in slow-roll, it is crucial to divide the inflationary epoch into three different regimes, which reflects the change of behaviour for a given mode when it crosses the Hubble scale. These regimes are conventionally called sub-horizon, horizon-crossing (transition), and super-horizon. Within all three regions analytic solutions for the perturbations valid to first order could be found. The sub-horizon region is irrelevant for the correlator of the gravitational potential. Relating the transition and super-horizon regions is not trivial, as there is no analytic result that is valid to first order at the boundary between them.

Using the procedure where we identify leading order asymptotic expansions, we could determine the relative normalization of the super-horizon solution with respect to the solution in the sub-horizon region using analytic properties.

To determine the solution for the scalar perturbations other than the gravitational potential in the super-horizon region we need a final cornerstone: the application of slow roll to the perturbations. For this it was essential that we treated the background using an arbitrary time variable, since the perturbed metric has to be rewritten in terms of a changed background metric. In particular this method was used to obtain an integral expression for the particular solution of the gravitational potential in terms of background quantities only. Although this expression is a priori not expected to be good near the end of (slow-roll) inflation, we show that it can actually be a very good approximation if $\tilde{\eta}^{\perp}$ goes to zero at the end of inflation.

Making some assumptions about the evolution of the universe and its matter content after inflation we gave first-order expressions for the adiabatic, isocurvature and mixing correlators of the gravitational potential at the time of recombination. Here we neglected the possible effects of (p)reheating, which are still under investigation. The adiabatic correlator includes the effect of entropy perturbations acting as a source for the gravitational potential during inflation and can be given in terms of background quantities expressed during the transition region (apart from possible end-of-inflation effects in the particular solution which are absent if $\tilde{\eta}^{\perp}$ goes to zero). The isocurvature correlator depends strongly on the transition at the end of inflation.

Finally, we discussed the example of multiple scalar fields on a flat manifold with a quadratic potential. To first order the trajectory of all fields through field space can be found in terms of one function of time, and the particular solution can be determined completely analytically using the slow-roll approximation on the perturbations. We concluded with an explicit numerical check of the amplitude of the adiabatic correlator and found this to be consistent with our analytical results.

Multiple-field effects are important in the adiabatic correlator of the gravitational potential if $\tilde{\eta}^{\perp}$ is sizable during the last 60 e-folds of inflation. The most important source of multiple-field effects is the particular solution of the gravitational potential. We found in our numerical example that this term can contribute even at leading order. Hence it can be dangerous to neglect this term, even when looking only at leading order. This contribution is included implicitly in the function $N(\boldsymbol{\phi})$ of [21], but we derived an explicit expression. If $\tilde{\eta}^{\perp}$ peaks in the transition region, the rotation of the basis induced by the background field dynamics over the transition region can be another source of multiple-field effects, but in generic situations we found it to be beyond the level of first order in slow roll. Although the details of the isocurvature and mixing correlators of the gravitational potential depend on the transition at the end of inflation, an important role is again played by the slow-roll function $\tilde{\eta}^{\perp}$. If it is zero there will be no mixing between adiabatic and isocurvature perturbations. Moreover, we found that with the assumption of an immediate transition to a radiation dominated universe at the end of inflation, only the e_2 component of q (using our basis) enters into the final expression for the isocurvature correlator, independently of the actual number of entropy perturbations.

Acknowledgments

This work is supported by the European Commission RTN programme HPRN-CT-2000-00131. S.G.N. is also supported by piority grant 1096 of the Deutsche Forschungsgemeinschaft and European Commission RTN programme HPRN-CT-2000-00148/00152.

Appendix A. Geometrical concepts

Consider a real manifold $\mathcal{M}$ with metric $\mathbf{G}$ and local coordinates $\boldsymbol{\phi} = (\phi^a)$. From the components of this metric G_{ab} the metric-connection Γ^a_{bc} is obtained using the metric postulate. The curvature tensor of the manifold can be introduced using tangent vectors $\boldsymbol{B}, \boldsymbol{C}, \boldsymbol{D}$:

$$[\mathbf{R}(\boldsymbol{B}, \boldsymbol{C})\boldsymbol{D}]^a \equiv R^a_{\ bcd} B^b C^c D^d \equiv \left(\Gamma^a_{bd,c} - \Gamma^a_{bc,d} + \Gamma^e_{bd}\Gamma^a_{ce} - \Gamma^e_{bc}\Gamma^a_{de}\right) B^b C^c D^d. \tag{A.1}$$

One should realize that for notational convenience we do not use the standard definition as made for example in [56]: our $\mathbf{R}(\boldsymbol{B}, \boldsymbol{C})\boldsymbol{D}$ is conventionally denoted by $\boldsymbol{R}(\boldsymbol{C}, \boldsymbol{D}, \boldsymbol{B})$.

The metric $\mathbf{G}$ introduces an inner product and the corresponding norm on the tangent bundle of the manifold:

$$\boldsymbol{A} \cdot \boldsymbol{B} = \boldsymbol{A}^\dagger \boldsymbol{B} \equiv \boldsymbol{A}^T \mathbf{G} \boldsymbol{B} = A^a G_{ab} B^b, \qquad |\boldsymbol{A}| \equiv \sqrt{\boldsymbol{A} \cdot \boldsymbol{A}}, \tag{A.2}$$

for any two vector fields $\boldsymbol{A}$ and $\boldsymbol{B}$. The cotangent vector $\boldsymbol{A}^\dagger$ is defined by $(\boldsymbol{A}^\dagger)_a \equiv A^b G_{ba}$. The Hermitean conjugate $\mathbf{L}^\dagger$ of a linear operator $\mathbf{L} : T_p\mathcal{M} \longrightarrow T_p\mathcal{M}$ with respect to this inner product is defined by

$$\boldsymbol{B} \cdot (\mathbf{L}^\dagger \boldsymbol{A}) \equiv (\mathbf{L}\boldsymbol{B}) \cdot \boldsymbol{A}, \tag{A.3}$$

so that $\mathbf{L}^\dagger = \mathbf{G}^{-1}\mathbf{L}^T\mathbf{G}$. A Hermitean operator $\mathbf{H}$ satisfies $\mathbf{H}^\dagger = \mathbf{H}$. An important example of Hermitean operators are the projection operators. Apart from being Hermitean, a projection operator $\mathbf{P}$ is idempotent: $\mathbf{P}^2 = \mathbf{P}$.

To complete our discussion on the geometry of $\mathcal{M}$ we introduce different types of derivatives. In the first place we have the covariant derivative on the manifold, denoted by ∇_a, which acts in the usual way, i.e.

$$\nabla_b A^a \equiv A^a_{\ ,b} + \Gamma^a_{bc} A^c \tag{A.4}$$

on a vector A^a. On a scalar function V, the derivative $\boldsymbol{\partial}$ and the covariant derivative $\boldsymbol{\nabla}$ are equal $(\boldsymbol{\nabla} V)_a = (\boldsymbol{\partial} V)_a \equiv V_{,a}$. If we represent $\mathrm{d}\boldsymbol{\phi}$ as a standing vector, $\boldsymbol{\nabla}$ and $\boldsymbol{\partial}$ are naturally lying vectors and therefore $\boldsymbol{\nabla}^T$ and $\boldsymbol{\partial}^T$ are standing vectors. The second covariant derivative of a scalar function V is a matrix with two lower indices: $(\boldsymbol{\nabla}^T \boldsymbol{\nabla} V)_{ab} = \nabla_a \nabla_b V$. The covariant derivative $\mathcal{D}_\mu$ with respect to the spacetime variable x^μ on a vector $\boldsymbol{A}$ of the tangent bundle is defined in components as

$$\mathcal{D}_\mu A^a \equiv \partial_\mu A^a + \Gamma^a_{bc} \partial_\mu \phi^b A^c, \tag{A.5}$$

while $\mathcal{D}_\mu$ acting on a scalar is simply equal to ∂_μ. Notice that the spacetime derivative of the background field $\partial_\mu\boldsymbol{\phi}$ and the field perturbation $\boldsymbol{\delta\phi}$ transform as vectors, even though the fields $\boldsymbol{\phi}$ in general do not, as they are coordinates on a manifold.

References

[1] Guth A H 1981 *Phys. Rev.* D **23** 347
[2] Linde A D 1990 *Particle Physics and Inflationary Cosmology* (Chur: Harwood Academic Publishers)
[3] Kinney W H, Melchiorri A and Riotto A 2001 *Phys. Rev.* D **63** 023505 (*Preprint* astro-ph/0007375)
[4] Hannestad S, Hansen S H, Villante F L and Hamilton A J S 2001 Constraints on inflation from CMB and Lyman-α forest *Preprint* astro-ph/0103047
[5] Bardeen J M, Steinhardt P J and Turner M S 1983 *Phys. Rev.* D **28** 679
[6] Mukhanov V F, Feldman H A and Brandenberger R H 1992 *Phys. Rep.* **215** 203
[7] Liddle A R and Lyth D H 1993 *Phys. Rep.* **231** 1 (*Preprint* astro-ph/9303019)
[8] Stewart E D and Lyth D H 1993 *Phys. Lett.* B **302** 171 (*Preprint* gr-qc/9302019)
[9] Martin J and Schwarz D J 2000 *Phys. Rev.* D **62** 103520 (*Preprint* astro-ph/9911225)
[10] Stewart E D and Gong J O 2001 *Phys. Lett.* B **510** 1 (*Preprint* astro-ph/0101225)
[11] Schwarz D J, Terrero-Escalante C A and Garcia A A 2001 *Phys. Lett.* B **517** 243 (*Preprint* astro-ph/0106020)
[12] Kodama H and Sasaki M 1984 *Prog. Theor. Phys. Suppl.* **78** 1
[13] Starobinsky A A 1985 *JETP Lett.* **42** 152
[14] Polarski D and Starobinsky A A 1992 *Nucl. Phys.* B **385** 623
[15] Garcia-Bellido J and Wands D 1996 *Phys. Rev.* D **53** 5437 (*Preprint* astro-ph/9511029)
[16] Garcia-Bellido J, Linde A D and Wands D 1996 *Phys. Rev.* D **54** 6040 (*Preprint* astro-ph/9605094)
[17] Nambu Y and Taruya A 1998 *Class. Quantum Grav.* **15** 2761 (*Preprint* gr-qc/9801021)
[18] Gordon C, Wands D, Bassett B A and Maartens R 2001 *Phys. Rev.* D **63** 023506 (*Preprint* astro-ph/0009131)
[19] Lyth D H and Riotto A 1999 *Phys. Rep.* **314** 1 (*Preprint* hep-ph/9807278)
[20] Sasaki M and Stewart E D 1996 *Prog. Theor. Phys.* **95** 71 (*Preprint* astro-ph/9507001)
[21] Nakamura T T and Stewart E D 1996 *Phys. Lett.* B **381** 413 (*Preprint* astro-ph/9604103)
[22] Mukhanov V F and Steinhardt P J 1998 *Phys. Lett.* B **422** 52 (*Preprint* astro-ph/9710038)
[23] Polarski D and Starobinsky A A 1994 *Phys. Rev.* D **50** 6123 (*Preprint* astro-ph/9404061)
[24] Langlois D 1999 *Phys. Rev.* D **59** 123512 (*Preprint* astro-ph/9906080)
[25] Hwang J-C and Noh H 2000 *Phys. Lett.* B **495** 277 (*Preprint* astro-ph/0009268)
[26] Hwang J-C and Noh H 2001 Cosmological Perturbations with Multiple Fluids and Fields *Preprint* astro-ph/0103244
[27] Starobinsky A A and Yokoyama J 1995 *Proc. 4th Workshop on General Relativity and Gravitation (Kyoto)* ed Nakao K *et al* (Kyoto: Kyoto University) p 381 (*Preprint* gr-qc/9502002)
[28] Hwang J-C and Noh H 2001 Gauge-ready formulation of the cosmological kinetic theory in generalized gravity theories *Preprint* astro-ph/0102005
[29] Starobinsky A A, Tsujikawa S and Yokoyama J 2001 *Nucl. Phys.* B **610** 383 (*Preprint* astro-ph/0107555)
[30] Linde A D 1994 *Phys. Rev.* D **49** 748 (*Preprint* astro-ph/9307002)
[31] Cremmer E, Ferrara S, Girardello L and Van Proeyen A 1983 *Nucl. Phys.* B **212** 413
[32] Nilles H P 1984 *Phys. Rep.* **110** 1
[33] Witten E 1985 *Phys. Lett.* B **155** 151
[34] Lahanas A B and Nanopoulos D V 1987 *Phys. Rep.* **145** 1
[35] Ferrara S, Lust D, Shapere A D and Theisen S 1989 *Phys. Lett.* B **225** 363
[36] Quevedo F 1996 Lectures on superstring phenomenology *Preprint* hep-th/9603074

[37] Groot Nibbelink S and Van Tent B J W 2000 Density perturbations arising from multiple field slow-roll inflation *Preprint* hep-ph/0011325

[38] Liddle A R, Parsons P and Barrow J D 1994 *Phys. Rev.* D **50** 7222 (*Preprint* astro-ph/9408015)

[39] Bardeen J M 1980 *Phys. Rev.* D **22** 1882

[40] Sasaki M 1986 *Prog. Theor. Phys.* **76** 1036

[41] Mukhanov V F 1988 *Sov. Phys. JETP* **67** 1297

[42] Mukhanov V F 1985 *JETP Lett.* **41** 493

[43] Martin J and Brandenberger R H 2001 *Phys. Rev.* D **63** 123501 (*Preprint* hep-th/0005209)

[44] Niemeyer J C 2001 *Phys. Rev.* D **63** 123502 (*Preprint* astro-ph/0005533)

[45] Lesgourgues J, Polarski D and Starobinsky A A 1997 *Nucl. Phys.* B **497** 479 (*Preprint* gr-qc/9611019)

[46] Martin J, Riazuelo A and Sakellariadou M 2000 *Phys. Rev.* D **61** 083518 (*Preprint* astro-ph/9904167)

[47] Taruya A and Nambu Y 1998 *Phys. Lett.* B **428** 37 (*Preprint* gr-qc/9709035)

[48] Sasaki M and Tanaka T 1998 *Prog. Theor. Phys.* **99** 763 (*Preprint* gr-qc/9801017)

[49] Kodama H and Sasaki M 1986 *Int. J. Mod. Phys.* A **1** 265

[50] Bucher M, Moodley K and Turok N 2000 *Phys. Rev.* D **62** 083508 (*Preprint* astro-ph/9904231)

[51] Bartolo N, Matarrese S and Riotto A 2001 *Phys. Rev.* D **64** 123504 (*Preprint* astro-ph/0107502)

[52] Wands D, Malik K A, Lyth D H and Liddle A R 2000 *Phys. Rev.* D **62** 043527 (*Preprint* astro-ph/0003278)

[53] Finelli F and Brandenberger R 2000 *Phys. Rev.* D **62** 083502 (*Preprint* hep-ph/0003172)

[54] Henriques A B and Moorhouse R G 2001 Cosmic microwave background and parametric resonance in reheating *Preprint* hep-ph/0109218

[55] Liddle A R, Mazumdar A and Schunck F E 1998 *Phys. Rev.* D **58** 061301 (*Preprint* astro-ph/9804177)

[56] Nakahara M 1996 *Geometry, Topology and Physics* (Bristol: Institute of Physics Publishing)